

Physicochemical and Nutritional Quality of Some Local and Modern Aromatic Rice Varieties of Bangladesh

Md. Farid Hossain*, Md. Sultan Uddin Bhuiya² and Masood Ahmed²

Department of Agronomy, Hajee Mohammad Danesh Science and Technology University, Dinajpur, Bangladesh ¹, Department of Agronomy, Bangladesh Agricultural University, Mymensingh, Bangladesh ²

Abstract

Physicochemical properties of five local and three modern aromatic rice varieties were studied. The parameters used for this study were milling outturn, head rice outturn, grain length, length breadth ratio, 1000 grain weight, amylose content, protein content, cooking time, grain elongation ratio, volume expansion ratio, gelatinization temperature, appearance and aroma. The parameters were found highly significant. Milling outturn of the studied varieties ranged from 69 to 71% and head rice outturn from 58 to 68%. All the rice varieties had translucent kernels and good in appearance. Eight samples had either medium slender, medium bold or short bold grains. The varieties had 6.1-6.9% protein and 23.6-25.5 % amylose content. Chinigura and BRRI dhan34 had highest protein and amylose content respectively. 1000 grain weight varied from 7.9 to 14.2 g. The cooking time of the varieties varied from 11.5 to 19.0 minutes. Kalizera and Badshabhog were more aromatic than other varieties. Elongation ratio and volume expansion ratio varied from 1.8 to 2.3 and from 3.5 to 4.6 respectively. Gelatinization temperatures of all varieties were low to intermediate.

Key words: Physicochemical properties, nutritional quality, aromatic rice

Introduction

Rice is the staple food of Bangladesh. Rice supplies more than 80% of the calories and about 50% of the protein in the diet of the general people of Bangladesh¹. It is also a source of some B vitamins. Along with nutritional qualities, the physical, cooking and eating qualities of rice are also important. Grain quality of rice is getting prominence in breeding programmes in the countries self sufficient in its production. Aromatic rice constitutes a small but an important sub- group of rice. These are rated best in quality and fetch much higher price than non-aromatic rice in the domestic and international market. Consumers' preference largely determines the commercial success of a rice variety.

Bangladesh Journal of Nutrition. Vol. 20-21, December 2008. Institute of Nutrition and Food Science, University of Dhaka-1000, Bangladesh.

* Author for correspondence

Preference for grain size and shape vary from one group of consumers to another². The rice millers prefer varieties with high milling and head rice outturn, whereas consumers consider quality³. The amylose content of rice is considered as the main parameter of cooking and eating quality⁴. Cooking behavior is one of the important determinants of quality⁵. The aroma of rice was significantly influenced by variety⁶. However; such rice should possess excellent physical and chemical characteristics acceptable to the local consumers and those abroad. The objective of this study was to determine the physicochemical properties associated with rice quality of five local and three modern aromatic rice varieties.

Materials and Methods

Five local and three modern aromatic rice varieties were grown in Hajee Mohammad Danesh Science and Technology University Farm, Dinajpur, Bangladesh during aman season of 2004 with three replications in randomized complete block design. They were milled raw and analyzed for physicochemical properties. Grain physicochemical parameters were measured at Grain Quality and Nutrition Division Laboratory, Bangladesh Rice Research Institute, Gazipur. Milled rice outturn was determined by dehulling 200g rough rice in a Satake Rice Mill, followed by 75 second polishing in a Satake Grain Testing Mill TM-05. Head rice outturn was determined by separating broken from milled rice by hand. Milled rice outturn and head rice outturn were expressed as percent of rough and milled rice respectively. Grain length and breadth were measured by slide calipers. In determining the size and shape, milled rice was first classified into three classes based on length, long (more than 6mm in length), medium(5-6mm in length) and short (less than 5mm in length).Then again classified into three classes according to the ratio of length to breadth; slender (ratio more than 3); bold (ratio2-3); round (ratio less than 2). Amylose content was determined by the procedure of Juliano⁷. Protein content was calculated from nitrogen and it was determined by the micro Kjeldahl method⁸. Appearance of the grain mainly depends on the amount of chalkiness, size, shape and colour of the grain. Aroma of cooked kernels was tested by the procedure of Sood and Siddiq⁹.Gelatinization temperature was determined according to the procedure of Little *et al*¹⁰. Volumes of cooked and milled rice were measured by water displacement. Data presented in the tables are mean of three replications.

Results and Discussion

Milling outturn of the studied samples varied from 69 to 71%. Most of the varieties had more than 70%. The head rice outturn varied between 58% and 68%. Out of eight varieties, 6 varieties had more than 66 % head rice outturn but long grain modern varieties BRRI dhan37, BRRI dhan38 had 58% and 59% head rice outturn respectively (Table 1). Biswas *et al*¹¹ studied 34 rice genotypes and found that milling outturn (%) ranged from 68 to 72 %.The head rice outturn was the proportion of the whole grain in

milled rice. It depends on the varietal character as well as drying condition¹². Translucent grains are attractive and fetch high market price. All the local and modern tested aromatic rice varieties had translucent kernels and good in appearance. Length and length breadth ratio of the reported varieties were 3.7 to 5.6 mm and 2.0 to 3.1 respectively (Table 1).

Table1. Physical properties of five local and three modern aromatic rice varieties

Variety	Milling outturn (%)	Head rice outturn (%)	1000 grain wt.(g)	Grain length (mm)	Length: breadth	*Size & shape	Appearance
Local							
Kataribhog	71	67	11.1	5.2	2.8	MB	Good
Radhunipagal	71	66	9.0	3.8	2.0	SB	Good
Chinigura	70	66	8.1	3.7	2.0	SB	Good
Badshabhog	71	68	7.9	3.7	2.2	SB	Good
Kalizera	71	66	8.8	3.8	2.1	SB	Good
Modern							
BRRi dhan34	71	67	8.4	3.8	2.1	SB	Good
BRRi dhan37	70	58	13.9	5.5	3.1	MS	Good
BRRi dhan38	69	59	14.2	5.6	3.1	MS	Good

*MS= Medium slender; MB= Medium bold; SB= short bold

Length of BRRi dhan37 (5.5mm) and BRRi dhan38 (5.6mm) were similar which were significantly higher than that of Kataribhog (5.2mm), Radhunipagal (3.8mm), Chinigura (3.7mm), Badshabhog (3.7mm), Kalizera (3.8mm) and BRRi dhan34 (3.8mm). Length breadth ratio of BRRi dhan37 (3.1) and BRRi dhan38 (3.1) were same, which was significantly higher than other varieties (Table1). Eight samples had either medium slender, medium bold or short bold. BRRi dhan37 and BRRi dhan38 had medium slender grains. On the other hand, Kataribhog had medium bold but Radhunipagal, Chinigura, Badshabhog, Kalizera and BRRi dhan34 were short bold. Amylose content influences the quality of cooked rice. High amylose rice varieties have high volume expansion and fluffy cooked rice. Amylose content of the tested varieties varied between 23.6% and 25.5% (Table 2). All the varieties were of intermediate amylose content (20-25%) except BRRi dhan34, which contained high amylose (more than 25 %). Intermediate amylose rice is preferred types in most of the rice growing areas of the world. Protein content of rice, which does not have much

Table 2. Protein content, amylose content and some cooking properties of five local and three modern aromatic rice varieties

Variety	Protein (%)	Amylose (%)	*Aroma	Cooking time (min.)	Elongation ratio	Volume expansion ratio	Gelatinization temperature
Local							
Kataribhog	6.3	24.7	++	15.0	1.8	4.0	Low
Radhunipagol	6.7	24.8	++	14.0	1.9	3.5	Low
Chinigura	6.9	23.6	++	12.5	2.0	3.7	Intermediate
Badshabhog	6.8	23.7	+++	11.5	2.0	3.7	Low
Kalizera	6.5	24.8	+++	12.0	2.0	3.7	Intermediate
Modern							
BRR1 dhan34	6.1	25.5	++	13.0	2.3	3.9	Low
BRR1 dhan37	6.2	23.8	+	19.0	2.0	4.6	Low
BRR1 dhan38	6.1	23.8	++	19.0	2.0	4.4	Low

* Aroma: Mild aromatic (+), moderately aromatic (++), strongly aromatic (+++)

influence on cooking properties, is important from nutritional point of view only. Protein content of the varieties varied from 6.1% to 6.9 % in brown rice (Table 2). The cooking time of these tested varieties varied between 11 and 19 minutes. BRR1 dhan37 and BRR1 dhan38 required maximum cooking time. Elongation ratio varied from 1.8 to 2.3 and volume expansion ranged from 3.5 to 4.6 (Table 2). BRR1¹³ observed that among 12 scented rice varieties amylose content varied from 22.0 to 25.9%, protein content varied from 6.5 to 9.0%, length of grain varied from 3.4 to 5.1 mm, length to breadth ratio varied from 1.9 to 3.4, cooking time varied from 11 to 14 minutes, elongation ratio ranged from 1.3 to 1.5 and volume expansion ratio ranged from 3.2 to 3.4. Gelatinization temperatures of all tested varieties were low to intermediate (Table 2). Dutt *et al*¹⁴ reported that Kataribkog was with less aroma in contrast to Kalizera, Badshabhog with intense aroma.

Acknowledgement

The authors wish to thank scientists of Grain Quality and Nutrition Division, Bangladesh Rice Research Institute (BRR1) for technical assistance throughout the study period.

References

1. Yusuf HKM. In: Report of the sustainable Food Security Mission in Bangladesh (FAO, Rome). Dhaka. 1997.
2. Khush GS, Paule CM and Dela Cruz NM. Rice grain quality evaluation and improvement at IRRI In: Proceedings of the workshop on chemical aspects of rice grain quality. IRRI, Los Banos, Philippines. 1979; pp21-31.

- 3 Merca FE and Juliano BO. Physicochemical properties of starch of intermediate amylose and starch. 1981; 33 (8):253-260.
- 4 Juliano BO. Physicochemical properties of starch and protein in relation to grain quality and nutritional value of rice. In: IRRI Rice Breeding. Los Banos Philippines. 1972; pp 389-405.
- 5 Feillet P and Marie R. Rice breeding for grain quality in France. In: Proceedings of the workshop on chemical aspects of grain quality. IRRI, Los Banos, Philippines. 1979; pp123-127.
- 6 Sikder MSI, Ahmed M and Hossain SMA. Effect of spacing and nitrogen fertilizer level on the yield and quality of some varieties of aromatic rice. M.S.Thesis in Agronomy. Dept. Agron. Bangladesh Agril. Univ, Mymensingh, Bangladesh. 2001.
- 7 Juliano BO. A simplified assay for milled rice amylose. Cereal Sci. Today. 1971; 16:334-338,340,360.
- 8 AOAC (Association of Official Agricultural Chemists). Methods of Analysis. 11th ed., Washington D.C. 1970; p 858.
- 9 Sood BC and Siddiq EA. A rapid technique for scent determination in rice. Indian J.Genet. Plant Breed. 1978; 38:268-71.
- 10 Little RR, Hilder GB and Dawson EH. Differential effect of dilute alkali on 25 varieties of milled white rice. Cereal Chem. 1958; 35:111-126.
- 11 Biswas SK, Banu B, Kabir KA, Begum F and Choudhury NH. Physicochemical properties of modern and local rice varieties of Bangladesh. Bangladesh Rice J 1992 ;(1&2):128-131.
- 12 Adair CR, Bollich CN, Bowman DH, Jodon NE, Johnston TH, Webb BD and Atkins JG. Rice breeding and testing methods in the United States. In: Rice in The United States: Varieties and Production. US Dept. Agr. Handbook 289 (Revised). 1973; pp 22-75.
- 13 BRRI (Bangladesh Rice Research Institute). Annual Internal Review. Bangladesh Rice Research Institute, Gazipur, Bangladesh. 1992; pp.1-18.
- 14 Dutta RK, Lahiri BP and Mia MAB. Characterization of some aromatic and fine rice cultivars in relation to their physicochemical quality of grains. Indian J. Plant physiology. 1998; 3(1):61-64.