

Comparative Antioxidative and Antihyperglycemic Profiles of Pneumatophores of Two Mangrove Species *Avicennia alba* and *Sonneratia apetala*

Biswajit Biswas, Mimi Golder, Tannami Islam and Samir Kumar Sadhu

Pharmacy Discipline, Khulna University, Khulna-9208, Bangladesh

(Received: July 31, 2018; Accepted: September 10, 2018; Published (web): December 10, 2018)

ABSTRACT: Ethanolic extracts of pneumatophores of two mangrove species- *Avicennia alba* and *Sonneratia apetala* were studied *in vitro* for antioxidant capacity by measuring their ability to scavenge free radicals and determining total phenolic, flavonoid and tannin contents. *In vivo* measurement of antihyperglycemic activity of extracts was done by oral glucose tolerance test. In considering the antioxidant activity, *S. apetala* extract showed superior IC₅₀ (concentration of sample required to inhibit 50% of free radicals) value for scavenging DPPH radical (71.77 µg/ml), hydrogen peroxide radical (97.27 mg/l), hydroxyl radical (79.62 mg/l) and superoxide anion (108.89 mg/l). For *A. alba*, the values for the radical scavenging assays were much higher. In addition, total phenol, flavonoid and tannin content demonstrated by *S. apetala* were 204.03 mgGAE/g, 228.68 mgQE/g and 235.89 mgGAE/g whereas for *A. alba*, they were 65.52 mgGAE/g, 44 mgQE/g and 37.71 mgGAE/g, respectively. In oral glucose tolerance test, *S. apetala* reduced the blood glucose level to a higher extent than *A. alba*. So, *S. apetala* with higher amount of secondary metabolites (phenol, flavonoid, tannin) is a superior source of natural antioxidants and antihyperglycemics.

Key words: *Avicennia alba*, *Sonneratia apetala*, pneumatophore, antioxidant, radical scavenger, secondary metabolites, antihyperglycemic.

INTRODUCTION

Any molecule containing one or more unpaired electrons are free radicals. The radicals are constantly produced during the normal oxidation of foods and about 1-4% of oxygen taken up in the body is converted to free radicals.¹ The free radical chain reactions are usually produced in the electron transport chain in mitochondria, through liver mixed function oxidases, xanthine oxidase and aldehyde oxidase activity and in macrophage by nitric oxide synthase enzyme. Free radicals mediate lipid peroxidation which in turn stimulate oxidation of sulfhydryl groups, glycation of protein, inactivation of enzymes and fragmentation of DNA bases and deoxyribose leading to cytotoxicity and mutations.²

Although the radicals are important intermediates in natural processes, a variety of pathological conditions like cancer, cardiovascular, inflammatory and respiratory diseases, infertility, aging as well as Parkinson's disease, Alzheimer's disease, multiple sclerosis, liver cirrhosis, muscular dystrophy etc. are caused by them.^{3,4} In addition, radicals or reactive oxygen species (ROS) together with increased triglyceride and lipoprotein levels produced during hyperglycemia due to impaired glucose metabolism play the central role in diabetes complications during diabetes mellitus. Along with oxidative stress, impaired insulin action leading to increased glucose production is also responsible for diabetes.⁵ Therefore, now the search is directed for antioxidative phytochemicals with antihyperglycemic therapeutic effects.

Correspondence to: Samir Kumar Sadhu
E-mail: <sksadhu1969@yahoo.com>
<sksadhu1969@pharm.ku.ac.bd>

Dhaka Univ. J. Pharm. Sci. 17(2): 205-211, 2018 (December)

DOI: <http://dx.doi.org/10.3329/dujps.v17i2.39177>

The body has several mechanisms to counteract the oxidative free radical damage by producing antioxidants. Antioxidants are scavengers of free radicals relative to oxidizable substrate that significantly delay or reduce substrate oxidation. However, single antioxidant molecule reacts with single free radical and hence a constant need to replenish antioxidant resources is important.⁶

Mangrove plants are equipped with very efficient free radical scavenging system due to the hostile environmental conditions like water logging, high salinity, low oxygen, high wind and temperature etc.⁷ They possess a wide variety of phytochemicals such as phenolic compounds (e.g. phenolic acids, flavonoids, quinones, coumarins, lignans, tannins), nitrogen compounds (alkaloids, amines, betalains), vitamins, terpenoids (including carotenoids) and some other endogenous metabolites rich in antioxidant and hypoglycemic activity.^{8,9}

Sundarbans is the largest mangrove forest of the world which is on the verge of extinction due to global warming, deforestation, soil erosion and devastating natural calamities.¹⁰ So, determination of antioxidative and antihyperglycemic activities of pneumatophores of two mangrove species *A. alba* and *S. apetala* along with their total contents of secondary metabolites e.g. phenols, flavonoids and tannins is much demanding and significant.

MATERIALS AND METHODS

Chemicals. Analytical grade reagents such as 2, 2-diphenyl-1-picryl hydrazyl (DPPH) and Folin-Ciocalteu reagent along with Na₂CO₃, NaNO₂, AlCl₃, NaOH, H₂O₂, Na₂HPO₄ · 2H₂O, NaH₂PO₄ · 2H₂O, 2-deoxy-2-ribose, FeCl₃, EDTA, TCA, TBA, PMS, ascorbic acid, NBT, NADH were used.

Plant materials. Pneumatophores of the two species *Avicennia alba* (Family: Acanthaceae) and *Sonneratia apetala* (Family: Lythraceae) were collected from Sundarbans (Boidyamari, Mongla region) in June, 2016 and identified by the experts at Forestry and Wood Technology Discipline, Khulna University, Khulna, Bangladesh.

Extracts preparation. Cold extraction was carried out for 200 gm of each plant powder with 96% ethanol. The percent yields for *A. alba* and *S. apetala* were 2.97 and 4.27, respectively.

Antioxidant Assay

Qualitative antioxidant activity test. The test was done using pre-coated silica gel TLC plates and developed in polar, medium polar and non-polar solvent systems. Then 0.02% DPPH solution was applied and color change was noted.¹¹

DPPH radical scavenging activity test. DPPH radical scavenging activity of sample extracts was tested by modifications of reported microplate method to determine IC₅₀ values using DPPH.¹² Thermo Scientific Multiskan Ex microplate photometer was used to measure the absorbance at 517 nm. Antioxidant capacity was determined in terms of IC₅₀, calculated from the log concentration-percent inhibition curve.¹³

Hydrogen peroxide scavenging activity test. It was measured in accordance with the method of Keser.¹⁴ Antioxidant activity was determined in terms of IC₅₀, calculated from the log concentration-percent inhibition curve.

Hydroxyl radical scavenging activity test. The test was performed according to the method described by Elizabeth and Rao with slight modifications.¹⁵ The activity was measured by determining the competition between deoxyribose and sample extracts for hydroxyl radicals generated by Fe³⁺/ascorbate/EDTA/H₂O₂ system (the Fenton reaction).¹⁶

Superoxide radical scavenging activity test. Superoxide radical scavenging assay is based on the reduction of nitro blue tetrazolium (NBT) in presence of NADH and phenazoniumme thosulphate (PMS).¹⁷ The test was performed according to the method described by Nishikimi with slight modifications.¹⁸

Determination of total secondary metabolites

Total phenolic content determination. Total phenolic content (TPC) was determined by using Folin-Ciocalteu reagent with gallic acid as standard

and expressed as mg gallic acid equivalent (mgGAE)/gm dried plant extract.¹⁹

Total flavonoid content determination.

Aluminium chloride colorimetric assay was used for the determination of total flavonoid content (TFC) of the sample extracts and quercetin was used as standard.²⁰ TFC of extracts, measured from quercetin in standard curve was expressed as mg quercetin in equivalent (mgQE)/gm dried plant extract

Total tannin content determination. Total tannin content (TTC) of sample extracts was determined by Folin-Ciocalteu agent and expressed as mg gallic acid equivalent (mgGAE)/gm dried plant extract.²¹

Antihyperglycemic assay. One of the acceptable methods to evaluate the antihyperglycemic activity of plant extracts is oral glucose tolerance evaluation or oral glucose tolerance test (OGTT).²² In this test, glucose is given to mice orally and level of glucose in blood sample is measured afterward to determine how quickly it is cleared from blood. In the present study, antihyperglycemic activity of ethanolic extract of

pneumatophores of *A. alba* and *S. apetala* was screened and compared with the glibenclamide standard.

RESULTS AND DISCUSSION

In the DPPH assay on TLC plate, any electron or hydrogen donating molecule in mixture will react with and bleach DPPH. Thus, DPPH is reduced from purple to a light yellow component.⁸ *A. alba* and *S. apetala* extracts showed free radical scavenging property indicated by the presence of light yellow spot on the TLC plate. In DPPH radical scavenging activity test, purple colored DPPH has maximum absorption at 517 nm. Scavenging DPPH radical by antioxidants results in a decrease in absorption over time which is proportional to the concentration of radicals being scavenged.²³ Figure 1 shows the DPPH radical scavenging assay results where *S. apetala* extract revealed an IC₅₀ value of 71.77 µg/ml whereas for *A. alba* it was 1196.74 µg/ml. So, compared to the standard ascorbic acid (IC₅₀ 14.42 µg/ml), *S. apetala* has relatively high antioxidant property than *A. alba*.




Figure 1. Comparison of % inhibition of DPPH radical by ascorbic acid and sample extracts.

Hydrogen peroxide (H₂O₂) itself is not a free radical rather a weak oxidizing agent that rapidly crosses the cell membranes and reacts with Fe²⁺ and Cu²⁺ ions to form hydroxyl radicals inside the cell.¹⁶ In the hydrogen peroxide scavenging assay, *A. alba* scavenged H₂O₂ with an IC₅₀ value of 178.65 mg/l

whereas for *S. apetala*, it was 97.27 mg/l. Again, IC₅₀ value for ascorbic acid standard was 83.95 mg/l (Figure 2). Therefore, *S. apetala* showed higher scavenging ability than *A. alba*.

In hydroxyl radical scavenging activity test, hydroxyl radicals were produced by incubating

ferric-EDTA with ascorbic acid and H_2O_2 at pH 7.4. Then the radical was reacted with 2-deoxy-2-ribose to generate a MDA-like product that forms a pink chromogen upon heating with TBA at low pH. The sample extracts added to the reaction mixture exerted antioxidant activity by competing with deoxyribose for hydroxyl radicals and thus preventing the

reaction.¹⁶ *A. alba* extract displayed hydroxyl radical scavenging activity with an IC_{50} value of 132.43 mg/l, for *S. apetala* it was 79.62 mg/l and for ascorbic acid standard, it was 62.95 mg/l (Figure 3). The IC_{50} values indicate that *S. apetala* plant extract is a better hydroxyl radical scavenger than the *A. alba*.




Figure 2. Comparison of % inhibition of H_2O_2 by ascorbic acid and sample extracts.




Figure 3. Comparison of % inhibition of hydroxyl radical by ascorbic acid and sample extracts.




Figure 4. Comparison of % inhibition of superoxide radical by ascorbic acid and sample extracts.

In the superoxide radical scavenging activity test, superoxide anion ($O_2^{\cdot-}$) derived from dissolved PMS-NADH coupling reaction reduces NBT to a purple form and hence NBT is the probe to quantify its ($O_2^{\cdot-}$) concentration. The reduction of absorbance at 560 nm after incubation of tested samples indicated the consumption of superoxide anion in the reaction mixture.²³ Figure 4 shows that the IC_{50} value for *A. alba* extract is 257.04 mg/l and for *S. apetala* it is 108.89 mg/L compared to the ascorbic acid standard (IC_{50} 92.26 mg/l). So, *S. apetala* is a better scavenger of superoxide radical than *A. alba*.

Plant secondary metabolites like polyphenols, flavonoids and tannins were suggested to have evolved for protecting the plant from herbivores.²⁴ In human, many phenolics are responsible for reducing the risk of developing chronic diseases like cardiovascular disease, cancer, diabetes, etc. Tannins have marked anti-tumor, anti-viral, and anti-HIV activities along with inhibition of lipid peroxidation and plasmin, mediation of DNA nicking and amelioration of renal failure etc. On the other hand, flavonoids exhibit anti-spasmodic, anti-inflammatory, and anti-allergic activities as well as protective effects against vascular and hepatic disorders. By investigating the mechanism of action of these effects, antioxidant and free-radical scavenging actions were found responsible.²⁵

In previous studies, different polyphenols were isolated from different parts of *A. alba* and *S. apetala*. Three new naphthoquinones, named avicequinone-A, -B, -C and their analogues avicenol A, B, C were reported in *A. alba*.²⁶ On the other hand, *S. apetala* was screened for secondary metabolites and different polyphenols, anthocyanins,

flavonoid aglycones and flavonoid glycosides along with gallic acid, (+)-catechin, (-)-epicatechin, ellagic acid and quercetin were that exhibit strong antioxidant activity.²⁷ Here, total phenolic content of *A. alba* was 65.52 mg gallic acid equivalent (mgGAE)/g and for *S. apetala* it was 204.03 mgGAE/g. However, total flavonoid content was 44.0 mgQE/g for *A. alba* and 228.68 mgQE/g for *S. apetala*. As far as total tannin content was concerned, *S. apetala* dominated with a value of 235.89 mgGAE/g, whereas for *A. alba* it was 37.71 mgGAE/g.

In considering the antihyperglycemic effect, oral glucose tolerance evaluation is a standard test for measuring how quickly exogenous glucose can be cleared from blood. Glucose uptake of cells from the blood is controlled by insulin and impairment in glucose tolerance (i.e., higher time to clear certain amount of glucose) indicates problems with conservation of glucose homeostasis i.e. carbohydrate metabolism, insulin resistance, diabetes etc.²⁸ In evaluating the antihyperglycemic activity of the pneumatophores of plant extracts, *S. apetala* showed significant effect rather than *A. alba* at doses 250 and 500 mg/kg body weight at 60 and 120 min after the administration of glucose (Table 1). Possible mechanisms of antihyperglycemic activity for the plant extract include substantial free radical scavenging activity, increased insulin releasing activity, insulin mimetic activity, altered glucose utilization, regeneration of pancreatic Islets of Langerhans and transport of blood glucose to the peripheral tissue mediated by the presence of secondary metabolites such as triterpenes, steroids, flavonoids, alkaloids, phenolics etc.⁹

Table 1. Blood glucose level (mmol/l) of plant extracts for oral glucose tolerance evaluation.

Sample extract	DRSA (IC_{50} μ g/ml)	HPSA (IC_{50} mg/l)	HRSA (IC_{50} mg/l)	SRSA (IC_{50} mg/l)	TPC (mg GAE/g)	TFC(mg QE/g)	TTC(mg GAE/g)
<i>A. alba</i>	1196.74	178.65	132.43	257.04	65.52	44.0	37.71
<i>S. apetala</i>	71.77	97.27	79.62	108.89	204.03	228.68	235.89

Values are means \pm S.D., n = 5, p < 0.05 (*), p < 0.01 (**).

Table 2. IC₅₀ values of different scavenging assays and total content of secondary metabolites (phenolics, flavonoid and tannin) of sample extracts.

Treatment group	Blood glucose level (mmol/l)		
	At 0 min	At 60 min	At 120 min
Control	5.88 ± 0.15	8.23 ± 0.78	7.70 ± 0.40
Glibenclamide	5.77 ± 0.12	4.17 ± 1.01**	3.58 ± 0.69**
<i>A. alba</i> (250 mg)	5.4 ± 0.49*	8.33 ± 0.91	7.45 ± 0.59
<i>A. alba</i> (500 mg)	5.38 ± 0.37**	8.28 ± 1.48	7.42 ± 0.63
<i>S. apetala</i> (250 mg)	4.18 ± 0.37**	7.92 ± 0.31	5.15 ± 0.40**
<i>S. apetala</i> (500 mg)	4.22 ± 0.33**	7.17 ± 0.45**	5.12 ± 0.31**

DRSA (DPPH Radical Scavenging Activity), HPSA (Hydrogen Peroxide Scavenging Activity), HRSA (Hydroxyl Radical Scavenging Activity), SRSA (Superoxide Radical Scavenging Activity), TPC (Total Phenolic Content), TFC (Total Flavonoid Content), Total Tannin Content (TTC).

When all the calculated data for the evaluation of antioxidative activity was accumulated in a single table (Table 2), a definite linear correlation was found between free radical scavenging effect and amount of total secondary metabolites of the samples. In addition, substantially higher antihyperglycemic activity of *S. apetala* was also supported by its superior contents of phenol, flavonoid, tannin and other secondary metabolites. So, on the basis of these results, it could be concluded that *S. apetala* might be a prospective resource of antioxidative and antihyperglycemic constituents.

CONCLUSION

Based on the obtained data of present study, it can be concluded that ethanolic extract of *S. apetala* pneumatophore, which contains large amounts of phenolic compounds, flavonoids and tannins, exhibits a higher extent of antioxidative free radical scavenging and antihyperglycemic activity than *A. alba* which contains less amount of secondary metabolites. These *in vitro* and *in vivo* assays are indicators of the plant extracts as potential source of natural antioxidants and antihyperglycemics which might be essential in the prevention of various oxidative stresses, diabetes and related physiological disorders.

ACKNOWLEDGEMENTS

This project was funded by the Ministry of Education, Bangladesh in advanced research aid

program in education and partly by Khulna University Research Cell (KURC). The authors would also like to thank the authority of Pharmacy Discipline, Khulna University for whole hearted support to carry out this research project and the experts at Forestry and Wood Technology Discipline, Khulna University for identification of plant samples.

REFERENCES

1. Veerabathran, S.K., Janaky, S. and Sekar, T. 2011. Antioxidant status of leaves of *Caesalpinia bonduca*. *Int. J. Pharm. App.* **2**, 262-266.
2. Ara, N. and Nur, H. 2009. *In vitro* antioxidant activity of methanolic leaves and flowers extracts of *Lippia alba*. *Res. J. Medicine Med. Sci.* **4**, 107-110.
3. Sarma, A.D., Mallick, A.R. and Ghosh, A.K. 2010. Free radicals and their role in different clinical conditions: an overview. *Int. J. Pharm. Sci. Res.* **1**, 185-192.
4. Pham-Huy, L.A., He, H. and Pham-Huy, C. 2008. Free radicals, antioxidants in disease and health. *Int. J. Biomed. Sci.* **4**, 89-96.
5. Bonnefont-Rousselot, D. 2002. Glucose and reactive oxygen species. *Curr. Opin. Clin. Nutr. Metab. Care.* **5**, 561-568.
6. Pourmorad, F., Hosseinimehr, S.J. and Shahabimajid, N. 2006. Antioxidant activity, phenol and flavonoid contents of some selected Iranian medicinal plants. *Afr. J. Biotechnol.* **5**, 1142-1145.
7. Thatoi, H.N., Patra, J.K. and Das, S.K. 2014. Free radical scavenging and antioxidant potential of mangrove plants: A review. *Acta. Physiol. Plant.* **36**, 561-579.
8. Masokoab, P. and Eloff, J.N. 2007. Screening of twenty-four South African *Combretum* and six *Terminalia* species (Combretaceae) for antioxidant activities. *Afr. J. Tradit. Complement. Altern. Med.* **4**, 231-239.

9. Das, S.K., Samantaray, D., Patra, J.K., Samanta, L. and Thatoi, H. 2016. Antidiabetic potential of mangrove plants: A review. *Front. Life Sci.* **9**, 75-88.
10. Aziz, A. and Paul, A.R. 2015. Bangladesh Sundarbans: Present status of the environment and biota. *Diversity* **7**, 242-269.
11. Dwivedy, B.N., Dabral, P. and Kumar, R. 2014. Antioxidant activity of ethanolic extract of *Heritierafomes* leaves. *World J. Pharm. Res.* **6**, 1713-1723.
12. Yokozawa, T., Chen, C.P., Dong, E., Tanaka, T., Nonaka, G. and Nishioka, I. 1998. Study on the inhibitory effect of tannins and flavonoids against the 1,1-diphenyl-2-picrylhydrazyl radical. *Biochem. Pharmacol.* **56**, 213-222.
13. Sadhu, S.K., Okuyama, E., Fujimoto, H. and Ishibashi, M. 2003. Separation of *Leucasaspera*, a medicinal plant of Bangladesh, guided by prostaglandin inhibitory and antioxidant activities. *Chem. Pharm. Bull.* **51**, 595-598.
14. Keser, S., Celik, S., Turkoglu, S., Yilmaz, O. and Turkoglu, I. 2012. Hydrogen peroxide radical scavenging and total antioxidant activity of Hawthorn. *Chem. J.* **2**, 9-12.
15. Elizabeth, K. and Rao, M.N.A. 1989. Oxygen radical scavenging activity of curcumin. *Intl. J. Pharm.* **58**, 237-240.
16. Hazra, B., Biswas, S. and Mandal, N. 2008. Antioxidant and free radical scavenging activity of *Spondiaspinnata*. *B.M.C. Complement. Altern. Med.* **8**, 1-10.
17. Elizabeth, S.B. and Murugesan, K. 2012. Evaluation of antioxidant potential of *Morinda citrifolia* L. root extract. *Int. Res. J. Pharmaceuticals.* **02**, 106-111.
18. Nishikimi, M., Rao, N.A. and Yagi, K. 1972. The occurrence of superoxide anion in the reaction of reduced phenazinemethosulphate and molecular oxygen. *Biochem. Biophys. Res. Commun.* **46**, 849-854.
19. McDonald, S., Prenzler, P.D., Autolovich, M. and Robards, K. 2001. Phenolic content and antioxidant activity of olive extracts. *Food Chem.* **73**, 73-84.
20. Marinova, D., Ribarova, F. and Atanassova, M. 2005. Total phenolics and total flavonoids in Bulgarian fruits and vegetables. *J. Univ. Chem. Technol. Metallurgy* **40**, 255-260.
21. Tambe, V.D. and Bhambar, R.S. 2014. Estimation of total phenol, tannin, alkaloid and flavonoid in *Hibiscus tiliaceus* Linn. wood extracts. *J. Pharmacogn. Phytochem.* **2**, 41-47.
22. Bnouham, M., Merhfour, F.Z., Ziyyat, A., Mekhfi, H., Aziz, M. and Legssyer, A. 2003. Antihyperglycemic activity of the aqueous extract of *Urticadioica*. *Fitoterapia* **74**, 677-681.
23. Liang, N. and Kitts, D.D. 2014. Antioxidant property of coffee components: Assessment of methods that define mechanisms of action. *Molecules* **19**, 19180-19208.
24. Banerjee, D., Chakrabarti, S., Hazra, A.K., Banerjee, S., Ray, J. and Mukherjee, B. 2008. Antioxidant activity and total phenolics of some mangroves in Sundarbans. *Afr. J. Biotechnol.* **7**, 805-810.
25. Yokozawa, T., Chen, C.P., Dong, E., Tanaka, T., Nonaka, G.I. and Nishiok, I. 1998. Study on the inhibitory effect of tannins and flavonoids against the 1,1-diphenyl-2-picrylhydrazyl radical. *Biochem. Pharmacol.* **56**, 213-222.
26. Ito, C., Katsuno, S., Kondo, Y., Tan, H.T.W. and Furukaw, H. 2000. Chemical constituents of *Avicenniaalba*: Isolation and structural elucidation of new naphthoquinones and their analogues. *Chem. Pharm. Bull.* **48**, 339-343.
27. Hossain, S.J., Iftekharuzzaman, M., Haque, M.A., Saha, B., Moniruzzaman, M., Rahman, M.M. and Hossain, H. 2016. Nutrient compositions, antioxidant activity and common phenolics of *Sonneratia apetala* (Buch.-Ham.) fruit. *Int. J. Food Prop.* **19**, 1080-1092.
28. Joy, K. and Kuttan, R. 1999. Anti-diabetic activity of *Picrorrhiza kurroa* extract. *J. Ethnopharmacol.* **67**, 143-148.