

Preliminary Antimicrobial Activity and Cytotoxicity of *Brunfelsia latifolia*

Rokeya Begum^{1*}, Mohammad S. Rahman², A. M. Sarwaruddin Chowdhury¹
and Mohammad A. Rashid^{2,3}

¹Department of Applied Chemistry and Chemical Technology, University of Dhaka, Dhaka-1000, Bangladesh

²Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Dhaka, Dhaka-1000, Bangladesh

³Centre for Biomedical Research, University of Dhaka, Dhaka-1000, Bangladesh

The *n*-hexane and chloroform soluble portions of the methanolic extract of *Brunfelsia latifolia* were subjected to antimicrobial screening and brine shrimp lethality bioassay. The *n*-hexane soluble fraction exhibited moderate antibacterial activity, while the *n*-hexane and chloroform soluble materials demonstrated significant cytotoxicity with LC₅₀ of 4.22 and 4.10 µg/ml, respectively.

The plant, *B. latifolia* (Synonym: *B. bonodora*, *B. australis*; Bengali name: Shusoma), is an evergreen shrub belonging to the Solanaceae family. The plant grows well in the tropical areas of the world and it is found all over Bangladesh.¹⁻³ The species of this genus are reported to have analgesic, anti-inflammatory and CNS depressant activities.^{4,5} Fruits of *Brunfelsia* are toxic to canine and small children.⁶ Previous phytochemical investigations with *Brunfelsia* led to the isolation of coumarins and alkaloids.⁷

The aerial parts of *B. latifolia* were collected from Bolda garden, Dhaka in January 2005. It was identified at the Department of Botany, University of

Dhaka. The stem bark of the plant was separated from the stem and cut into small pieces, cleaned, dried and pulverized.

The powdered stem bark (200 g) of *B. latifolia* was soaked in 500 ml methanol, filtered and concentrated using a rotary evaporator at low temperature (36-40°C) and reduced pressure. An aliquot (5 g) of the concentrated methanol extract was fractionated by the modified Kupchan partitioning protocol⁸ into *n*-hexane, carbon tetrachloride and chloroform to afford *n*-hexane (1.7 g), carbon tetrachloride (0.4 g), chloroform (1.3 g) and 1.4 mg aqueous soluble materials (1.4 g).

The preliminary antimicrobial activity of the extractives was determined at 400 µg/disc by the disc diffusion method⁹ against a number of Gram positive and Gram negative bacteria and fungi (Table 1). The bacterial and fungal strains used in this experiment were collected as pure cultures from the Institute of Nutrition and Food Science (INFS), University of Dhaka. Here standard Kanamycin (30 µg/disc) was used as the reference.

For *in vivo* cytotoxicity screening, DMSO solutions of the plant extracts were applied against *Artemia salina* for 24 hours.¹⁰ For the experiment 4 mg of each of the plant extracts was dissolved in DMSO and by serial dilution technique, solutions of

Correspondence to: Mohammad A. Rashid
Tel.: 880-2-8612069, 9661900-73, extn.- 4363, 8137;
Fax: 880-2-8612069, E-mail address: rashidma@aitbd.net
^{*}On leave from Department of Agricultural Chemistry, Sher-e-Bangla Agricultural University, Dhaka-1207, Bangladesh

varying concentrations such as 400, 200, 100, 50, 25, 12.5, 6.25, 3.125, 1.563, 0.781 µg/ml were obtained. After 24 hours, the median lethal concentration, LC₅₀, of the test samples was obtained by a plot of percentage of shrimps killed against the logarithm of the sample concentration.

The *n*-hexane and chloroform soluble fractions of the methanolic extract of *B. latifolia* showed the average zone of inhibition of 10-15 mm and 7-14 mm, respectively at a concentration of 400 µg/disc. The *n*-hexane solubles of the bark strongly inhibited the growth of *B. cereus*, *S. bodii* and *C. albicans*

having the zone of inhibition of 15 mm. The growth of *P. aureus* (14 mm), *A. niger* (14 mm) and *S. lutea* (13 mm) was also inhibited. It also showed mild inhibition of growth of *E. coli*, *S. dysenteriae* and *S. cerevacee*. On the other hand, the chloroform soluble material demonstrated mild activity against most of the organisms. In case of fungal strains, the *n*-hexane soluble materials showed strong inhibitory activity against *C. albicans* (15 mm) and *A. niger* (15 mm). However, the chloroform extracts revealed mild inhibitory activity against the fungal growth.

Table 1. Preliminary antimicrobial activity of *B. latifolia* extractives.

Test bacteria and fungi	Diameter of zone of inhibition (mm)		
	HX (400 µg/disc)	CF (400 µg/disc)	KAN (30 µg/disc)
Gram positive bacteria			
<i>Bacillus cereus</i>	15	10	40
<i>B. megaterium</i>	--	14	50
<i>B. subtilis</i>	10	08	40
<i>Sarcina lutea</i>	13	07	24
<i>Staphylococcus aureus</i>	10	08	45
Gram negative bacteria			
<i>Escherichia coli</i>	12	10	26
<i>Pseudomonas aeruginosa</i>	14	08	23
<i>Salmonella typhi</i>	10	08	30
<i>Salmonella paratyphi</i>	10	07	50
<i>Shigella boydii</i>	15	08	23
<i>Shigella dysenteriae</i>	12	07	24
<i>Vibro mimicus</i>	10	07	45
<i>V. parahemolyticus</i>	10	08	30
Fungus			
<i>Aspergillus niger</i>	14	11	20
<i>Candida albicans</i>	15	08	25
<i>Sacharomyces cerevaceae</i>	12	10	23

HX: *n*-hexane soluble fraction of the methanol extract; CF: chloroform soluble fraction of the methanol extract; KAN: Kanamycin; "--": indicates no activity.

The *n*-hexane and chloroform soluble fractions of the methanol extract were subjected to brine shrimp lethality bioassay by using the method of Meyer.¹⁰ In the brine shrimp lethality bioassay, the LC₅₀ values exhibited by the *n*-hexane and chloroform soluble fractions were found to be 4.22 µg/ml and 4.10 µg/ml, respectively. The standard,

vincristine sulphate, showed the LC₅₀ value of 0.23 µg/ml. It is clearly evident from the experiment that the *n*-hexane and chloroform soluble fractions demonstrated strong cytotoxic activity. The antimicrobial activity and cytotoxicity revealed by the various extractives of *B. latifolia* substantiate the folk uses of this plant in various diseases.

REFERENCES

1. Khan, M.S. and Mia, C. 2002. *Flora of Bangladesh, Solanaceae*, (Khan, M.S. and Rahman, M.M. Eds.) Bangladesh National Herbarium, Ministry of Environment and Forest, Govt. of the Peoples Republic of Bangladesh, vol. **53**, pp. 5.
2. Usher, G. 1984. *A Dictionary of Plants*. CBS publishers and distributors, Delhi-110032 (India), pp. 103.
3. Baytop, A. 1978. *Solanaceae*. In: *Flora of Turkey and the East Aegean Islands*. P. H. Davis (ed.), University press, Edinburgh, Vol. **6**, pp. 437-458.
4. Iyer, R.P., Brown, J.K., Chaubal, M.G and Malone, M.H. 1977. *Brunfelsia hopeana* I: Hippocratic screening and antiinflammatory evaluation. *Lloydia* **40**, 356-360.
5. Ruppelt, B.M., Pereira, E.F., Gonçalves, L.C. and Pereira, N.A. 1991. Pharmacological screening of plants recommended by folk medicine as anti-snake venom--I. Analgesic and anti-inflammatory activities. *Mem. Inst. Oswaldo. Cruz.* **86**, 203-205.
6. Spainhour, C.B., Fiske, R.A., Flory, W. and Reagor, J.C. J. 1990. A toxicological investigation of the garden shrub *Brunfelsia calcyina* var. *floribunda* (yesterday-today-and-tomorrow) in three species. *Vet. Diagn. Invest.* **2**, 3-8.
7. *Dictionary of natural products*, Chapman and Hall, 2002.
8. Vanwagenen, B.C., Larsen, R., Cardellina, J.H., Ran dazzo, D., Lidert, Z.C. and Swithenbank, C. 1993. Ulosantoin, a potent insecticide from the sponge *Ulosa ruetzleri*. *J. Org. Chem.* **58**, 335-337.
9. Bauer, A.W., Kirby, W.M.M., Sherris, J.C. and Truck, M. 1966. Antibiotic susceptibility testing by standard single disc diffusion method. *Am. J. Clin. Pathol.* **45**, 426-493.
10. Meyer, B. N., Ferringni, N.R., Puam, J. E., Lacobsen, L. B., Nicols, D. E. and McLaughilin, J.L. 1982. Brine Shrimp: A convenient general bioassay for active constituents. *Planta Med.* **45**, 31-32.