

## Phytochemical and Biological Investigations of *Acokanthera spectabilis*

Tamanna Ferdous Shanta<sup>1</sup>, Mohammad S. Rahman<sup>2</sup>, Shaila Kabir<sup>2</sup>,  
A.N.M. Hamidul Kabir<sup>1</sup>, Bilkis Begum<sup>3</sup> and Mohammad A. Rashid<sup>2,4</sup>

<sup>1</sup>Department of Applied Chemistry and Chemical Technology, University of Dhaka, Dhaka-1000, Bangladesh

<sup>2</sup>Phytochemical Research Laboratory, Department of Pharmaceutical Chemistry, Faculty of Pharmacy,  
University of Dhaka, Dhaka-1000, Bangladesh

<sup>3</sup>Department of Clinical Pharmacy and Pharmacology, University of Dhaka, Dhaka-1000, Bangladesh

<sup>4</sup>Centre for Biomedical Research, University of Dhaka, Dhaka-1000, Bangladesh

The *n*-hexane, carbon tetrachloride and chloroform soluble fractions of a methanol extract of *Acokanthera spectabilis* were subjected to antimicrobial screening and brine shrimp lethality bioassay. All of the partitionates showed moderate to strong inhibitory activity to microbial growth, while the carbon tetrachloride soluble fraction demonstrated highest cytotoxicity having LC<sub>50</sub> 6.16 µg/ml.

*Acokanthera spectabilis* (Family- Apocynaceae) is a fragrant ornamental shrub. The plant has reputation for its cardiogenic activities.<sup>1</sup> Previous phytochemical investigation with *A. spectabilis* led to the isolation of acobioside A and 14-*O*-acetylacovenodose C.<sup>2</sup> In the present study, we report the presence of a cardenolide as well as antimicrobial activity and cytotoxicity of the extractives of *A. spectabilis* growing in Bangladesh.

The <sup>1</sup>H NMR spectra were recorded using a Bruker AMX-400 (400 MHz) instrument and the NMR spectra were acquired in CDCl<sub>3</sub> and the δ values for <sup>1</sup>H spectra were referenced to the residual nondeuterated solvent signal.

---

**Correspondence to:** Mohammad A. Rashid  
Tel.: 880-2-8612069, 9661900-73,  
Extn. - 4363, 4364, 8137; Fax: 880-2-8612069  
E-mail address: rashidma@aitfbd.net; rashid\_phdu@yahoo.com

The leaves and stem of *A. spectabilis* were collected from Baldah garden, Dhaka in the month of September 2005. A voucher specimen for this collection has been deposited in the herbarium of the Department of Botany, University of Dhaka.

Fresh leaves and stem of *A. spectabilis* were collected, dried and ground to a coarse powder. The powdered sample (149 g) was subjected to cold extraction with methanol for about 7 days and filtered through a cotton plug followed by Whatman filter paper number 1. The extract was then concentrated by using a rotary evaporator. A portion (5 mg) of the concentrated methanol extract was fractionated by the modified Kupchan partitioning method<sup>3</sup> into *n*-hexane, carbon tetrachloride, chloroform and aqueous soluble fractions. Evaporation of solvents afforded *n*-hexane (0.50 g), carbon tetrachloride (0.65 g), chloroform (1.30 g) and aqueous soluble (1.20 g) materials.

The carbon tetrachloride soluble fraction (300 mg) was fractionated over Sephadex LH-20. The column was eluted with *n*-hexane - dichloromethane - methanol (2:5:1) mixtures to provide 25 fractions, each 10 ml. Preparative thin layer chromatography (stationary phase- silica gel F<sub>254</sub>, mobile phase - 10 % ethylacetate in toluene, thickness of plates-0.5 mm) of fractions 5-8 afforded a pure compound.

The  $^1\text{H-NMR}$  spectrum (400 MHz,  $\text{CDCl}_3$ ) of the purified compound indicated the presence of a cardenolide type glycoside.<sup>8</sup> However, due to poor spectral resolution and insufficient data, the structure of the compound could not be established.

The antimicrobial activity of the Kupchan fractions was determined by the disc diffusion method.<sup>4</sup> The bacterial strains listed in Table-1 were collected as pure cultures from the Institute of Nutrition and Food Science (INFS), University of Dhaka, Bangladesh. The samples were dissolved separately in chloroform and applied to sterile filter paper discs at a concentration of 400 and 200  $\mu\text{g}/\text{disc}$  for crude extractives and isolated cardenolide, respectively. Discs were then carefully dried to evaporate the residual solvent. Standard kanamycin

(30  $\mu\text{g}/\text{disc}$ ) was used as the positive control in the experiment.

For cytotoxicity screening, DMSO solutions of the plant extractives were applied against *Artemia salina* in a 1-day *in vivo* assay, the experimental details of which could be found elsewhere.<sup>5-7</sup> For the experiment, 4 mg of each of the Kupchan fraction was dissolved in DMSO and solutions of varying concentrations (400, 200, 100, 50, 25, 12.50, 6.25, 3.125, 1.563, 0.781  $\mu\text{g}/\text{ml}$ ) were obtained by serial dilution technique.

For each of the extractives, three samples were prepared for each of the bioassay. The zone of inhibition and  $\text{LC}_{50}$  were calculated as mean  $\pm$  SD ( $n=3$ ) for the antimicrobial screening and brine shrimp lethality bioassay, respectively.

**Table 1. Antimicrobial activity of the extractives of *A. spectabilis***

Test microorganisms	Diameter of zone of inhibition (mm)				
	HX (400 $\mu\text{g}/\text{disc}$ )	CT (400 $\mu\text{g}/\text{disc}$ )	CF (400 $\mu\text{g}/\text{disc}$ )	C (400 $\mu\text{g}/\text{disc}$ )	Kan (200 $\mu\text{g}/\text{disc}$ )
<b>Gram positive bacteria</b>					
<i>Bacillus cereus</i>	08.30 $\pm$ 1.21	11.33 $\pm$ 0.33	09.21 $\pm$ 1.36	09.12 $\pm$ 1.33	26.13 $\pm$ 1.83
<i>Bacillus megaterium</i>	11.21 $\pm$ 1.32	10.25 $\pm$ 1.57	--	--	29.21 $\pm$ 0.87
<i>Bacillus subtilis</i>	--	16.35 $\pm$ 1.34	13.25 $\pm$ 1.37	--	23.33 $\pm$ 1.11
<i>Staphylococcus aureus</i>	10.14 $\pm$ 0.94	14.25 $\pm$ 1.34	12.22 $\pm$ 1.19	08.16 $\pm$ 1.38	25.33 $\pm$ 1.67
<i>Sarcina lutea</i>	--	12.34 $\pm$ 2.02	11.45 $\pm$ 1.26	10.28 $\pm$ 1.37	24.47 $\pm$ 0.33
<b>Gram negative bacteria</b>					
<i>Escherichia coli</i>	09.25 $\pm$ 1.33	13.46 $\pm$ 1.44	11.50 $\pm$ 1.65	--	25.33 $\pm$ 1.13
<i>Pseudomonas aeruginosa</i>	08.34 $\pm$ 0.97	11.36 $\pm$ 1.26	09.34 $\pm$ 1.74	--	25.21 $\pm$ 0.54
<i>Salmonella paratyphi</i>	08.16 $\pm$ 1.67	11.11 $\pm$ 1.54	--	09.30 $\pm$ 1.48	20.27 $\pm$ 0.33
<i>Salmonella typhi</i>	11.41 $\pm$ 2.34	13.32 $\pm$ 1.29	17.02 $\pm$ 1.37	09.29 $\pm$ 0.87	25.24 $\pm$ 0.42
<i>Shigella dysenteriae</i>	09.35 $\pm$ 1.26	15.43 $\pm$ 1.64	12.35 $\pm$ 1.09	-	24.18 $\pm$ 1.33
<i>Vibrio mimicus</i>	10.29 $\pm$ 1.57	12.26 $\pm$ 0.33	13.29 $\pm$ 1.31	11.23 $\pm$ 1.11	26.17 $\pm$ 1.12
<i>Vibrio parahemolyticus</i>	10.31 $\pm$ 1.24	10.16 $\pm$ 1.32	15.22 $\pm$ 0.92	09.33 $\pm$ 1.23	25.26 $\pm$ 0.61
<b>Fungi</b>					
<i>Candida albicans</i>	11.34 $\pm$ 1.67	16.05 $\pm$ 1.15	11.11 $\pm$ 0.67	--	26.28 $\pm$ 0.91
<i>Aspergillus niger</i>	08.27 $\pm$ 1.36	11.25 $\pm$ 1.47	--	--	27.37 $\pm$ 1.02
<i>Sacharomyces cerevaceae</i>	--	15.41 $\pm$ 1.32	15.41 $\pm$ 1.06	--	27.26 $\pm$ 0.33

The diameters of zones of inhibition are expressed as mean  $\pm$  SD ( $n=3$ ); a diameter less than 8 mm was considered inactive; HX: *n*-hexane soluble fraction of the methanolic extract; CT: carbon tetrachloride soluble fraction of the methanolic extract; CF: chloroform soluble fraction of the methanolic extract; C: cardenolide; Kan: standard kanamycin

In the antimicrobial screening, the zones of inhibition produced by the *n*-hexane, carbontetrachloride and chloroform soluble fractions were found to be 8-11 mm, 10-16 mm and 9-17 mm, respectively at a concentration of 400  $\mu\text{g}/\text{disc}$  (Table 1). The zones of inhibition produced by cardenolide at a dose level of 200  $\mu\text{g}/\text{disc}$  were 9-11 mm (Table 1).

Three crude extractives (*n*-hexane, carbon tetrachloride and chloroform soluble partitionates) were screened by brine shrimp lethality bioassay for probable cytotoxic activity. The  $\text{LC}_{50}$  values of *n*-hexane, carbontetrachloride and chloroform fractions were found to be 26.91  $\mu\text{g}/\text{ml}$ , 6.16  $\mu\text{g}/\text{ml}$  and 12.58  $\mu\text{g}/\text{ml}$  (Table 2), respectively. The positive

control, vincristine sulphate, showed  $LC_{50}$  of 0.32  $\mu\text{g/ml}$ . From the results of the brine shrimp lethality bioassay it can be well predicted that the crude extracts have considerable cytotoxic potency. It is evident that carbon tetrachloride soluble materials has significant cytotoxic principles from which the cardenolide analog was isolated. But cytotoxic activity of the pure compound could not be performed due to lack of the test sample. Further study is warranted to identify the bioactive principle from *A. spectabilis*.

**Table 2.**  $LC_{50}$  data of test samples of *A. spectabilis*

Samples	$LC_{50}$ ( $\mu\text{g/ml}$ )
Vincristine sulphate	$0.320 \pm 0.11$
HX	$26.91 \pm 1.21$
CT	$06.16 \pm 0.33$
CF	$12.58 \pm 2.10$

The values of  $LC_{50}$  are expressed as mean  $\pm$  SD (n=3). VS: vincristine sulphate (Std.); HX: *n*-hexane soluble fraction of the methanolic extract; CT: carbontetrachloride soluble fraction of the methanolic extract; CF: chloroform soluble fraction of the methanolic extract

## REFERENCES

1. Radford, D.J., Gillies, A.D., Hinds, J.A. and Duffy P. 1986. Naturally occurring cardiac glycosides. *Med. J. Aust.* **144**, 540-544.
2. Karawya, M.S., Abdel-Wahab, S.M. and Niazi, H.M. 1974. Assay of cardenolides in *Acokanthera spectabilis*. *Planta Med.* **25**, 17-21.
3. Vanwagenen, B.C., Larsen, R., Cardellina, J.H., Ran dazzo, D., Lidert, Z.C. and Swithenbank, C. 1993. Ulosantoin, a potent insecticide from the sponge *Ulosa ruetzleri*. *J. Org. Chem.* **58**, 335-337.
4. Bauer, A.W., Kirby, W.M.M., Sherris, J.C. and Truck, M. 1966. Antibiotic susceptibility testing by standard single disc diffusion method. *Am. J. Clin. Pathol.* **45**, 426-493.
5. Meyer, B.N., Ferringni, N.R., Puam, J.E., L Jacobsen, L.B., Nichols, D.E. and McLaughlin, J.L. 1982. Brine shrimp: a convenient general bioassay for active constituents. *Planta Med.* **45**, 31-32.
6. McLaughlin, J.L., Rogers, L.L. 1998. The use of biological assays to evaluate botanicals. *Drug Info. J.* **32**, 513-524.
7. Persoone, G. 1980. *Proceeding of the international symposium on brine shrimp, Artemia salina*, pp. 1-3, University Press.
8. Huq, M.M., Jabbar, A., Rashid, M.A. and Hasan, C.M. 1999. A novel antibacterial and cardiac steroid from the roots of *Nerium oleander*. *Fitoterapia* **70**, 5-9.