

# Selective Cytotoxicity Evaluation in Anticancer Drug Screening of *Boehmeria virgata* (Forst) Guill Leaves to Several Human Cell Lines: HeLa, WiDr, T47D and Vero

Wardihan<sup>1</sup>, Muhammad Rusdi<sup>2</sup>, Gemini Alam<sup>3</sup>, Lukman<sup>4</sup>  
and Marianti A. Manggau<sup>4</sup>

<sup>1</sup>Faculty of Medicine, Hasanuddin University 90245, East Sulawesi Makassar, Indonesia

<sup>2</sup>Faculty of Mathematics and Natural Sciences, Islamic University of Makassar 90245, Makassar, Indonesia

<sup>3</sup>Department of Pharmacognosy and Phytochemistry, Faculty of Pharmacy, Hasanuddin University 90245 Makassar, Indonesia

<sup>4</sup>Department of Biopharmacy, Faculty of Pharmacy, Hasanuddin University 90245, East Sulawesi Makassar Indonesia 90245, Makassar, Indonesia

**ABSTRACT :** Cancer is one of the leading causes of death worldwide. Many researchers have attempted to develop new treatments that will improve the prognosis of cancer patients. Indonesian forests, one of the most important world's biodiversity, but little is known about chemical and/or pharmacological potential of these plants. The aim of this study to investigate selective cytotoxicity of *B. virgata* against of various cancer cell lines. The ethanolic extract of *B. virgata* leaves can act against HeLa, WiDr, T47D and Vero cell lines with IC<sub>50</sub> 18.991 ± 0.234, 18.925 ± 1.277, 12.732 ± 0.945 and 16.022±0.663 µg/ml with selective index 0.844, 0.847, 1.258 and 1.000, respectively. We highlight the significant cytotoxic effect of *B. virgata* from the leaves, which introduces promising expectations for new projects in chemistry, pharmacology and toxicology although non selective in these cell lines.

**Key words:** *B. virgata*, selective cytotoxicity, MTT, HeLa, WiDr, T47D and Vero.

## INTRODUCTION

Cancer is a leading cause of disease worldwide and was estimated that 12.7 million new cancer cases occurred worldwide.<sup>1</sup> An estimated 12.66 million people were diagnosed with cancer was estimated to account for around 14% of all deaths (due to any cause) worldwide.<sup>2</sup>

Plants play an important role as a source of effective anticancer agents. Currently, over 60% of used anticancer agents are derived in one way or another from natural sources, including plants, marine animals and microorganisms.<sup>3</sup> There are worldwide efforts to discover new anticancer agents from plants.

Many of the drugs, which we use today are based on folk remedies and subsequent ethnopharmacological studies and the traditional medicines are generally more acceptable from a cultural and spiritual perspective.<sup>4</sup> The *B. virgata* leaf have been widely used in Traditional Makassar Medicine to treat cancer.<sup>5</sup> *B. virgata* is classified in the family Urticaceae.<sup>6</sup> The genus *Boehmeria* have been widely studied by several author, in search of answers to their cytotoxic effect<sup>7</sup> but no *B. virgata* was reported yet.

In this investigation, cytotoxic effects, IC<sub>50</sub> and selectivity of ethanolic extract of *B. virgata* leaves on cancer (HeLa, WiDr and T47D) and normal (Vero) cell lines were studied, using the MTT reduction test.

---

**Correspondence to:** Lukman  
Tel +62 8971561010,  
Email: lukman\_m01@yahoo.co.id

## MATERIAL AND METHODS

**Plant material.** Leaves of *B. virgate* were collected from Malino (South Sulawesi, Makassar, Indonesia) in the month of March, the plant was identified and authenticated. Fresh plant leaves were shade dried at room temperature and ground into a fine powder.

**Extraction.** Five hundred gram of plant materials was used for extraction using ethanol by maceration method. The extraction time was fixed for 24 h. The extract was concentrated by rotary evaporator and dried through lyophilization.<sup>8</sup>

**Cell cultures.** We used 3 cell lines: HeLa as servix cell line, WiDr as a colon adenocarcinoma, T47D as human ductal breast epithelial tumor cell line and Vero as normal cell line. HeLa and T47D were cultured in RPMI 1650, WiDr was cultured in Dulbecco's Modified Eagle Medium (DMEM). Vero was cultured in M199 medium. All cells were subcultured after mild trypsinization with trypsin-EDTA (Sigma-Aldrich, USA), trypan blue dye (Sigma-Aldrich, USA) exclusion assay was performed to determine the cell number and viability. All media (Sigma-Aldrich, USA) were supplemented at 10% with fetal bovine serum (Gibco) and streptomycin plus penicillin (100 ug/ml and 100 u/ml, respectively; Sigma-Aldrich, USA). The cell line were kept at 37°C, 98% relative humidity with 5% CO<sub>2</sub> atmosphere.

**Cytotoxic assay.** This assay was carried out for 24 h to evaluate the cytotoxicity effect.<sup>9</sup> Sufficient number of exponentially growing cells to avoid confluence of the culture during treatments were seeded at 10.000 cells/mL to evaluate possible cytotoxic effect in Iwaki 96-well plates. The treatment started 24 h after seeding (to improve environment adaptation), after 24 h of treatment, the medium was replaced and cultures were maintained for another 24 h before the cell viability quantification. Control was always treated with the same amount of DMSO as used in the corresponding experiments.

**Cell viability: MTT test.** The tetrazolium salt MTT was used as an indicator of mammalian cell

survival and proliferation.<sup>10</sup> Yellow 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium bromide (MTT) method consists in the absorption of yellow tetrazolim salts by mitochondrial reductases of metabolically active cells, called formazan. Formazan accumulated in intracellular cell, is extracted by adding an organic solvent.<sup>11</sup> Solubilised formazan reagent is measured by Elisa microplate reader at 595 nm.<sup>12</sup> The culture in 96-well plates were incubated with 100 µl of fresh supplemented medium and 100 µL of MTT (5 mg/ml). The MTT formazan that produced by the cultured cell was added 100 µl of DMSO to each well to solubilize the MTT formazan. After incubate for 24 h at room temperature, the plate were read with an Elisa Reader at 595 nm (bio-Rad).

Percentage inhibition was calculated as follows:

Percentage inhibition (%) =

$$\frac{\text{OD of control} - \text{OD of treatment}}{\text{OD of control}} \times 100$$

**Data analysis:** All the experiment were performed in triplicate at least. Dose response curves were plotted and the IC<sub>50</sub> values (concentrations at which cellular effects are inhibited by 50%) were calculated using linear regression analysis compared with untreated cells.

## RESULTS AND DISCUSSION

Cytotoxic effect of ethanol extract from *B. virgate* leaves was tested against HeLa using colorimetric method MTT assay. All the cells were exposed to various concentration: 1.95; 3.91; 7.81; 15.63; 31.25; 62.50 and 125.00 µg/ml (Figure 1).

Cytotoxic effect of ethanol extract from *B. virgate* leaves was tested against WiDr using colorimetric method MTT assay. All the cells were exposed to various concentration: 1.95; 3.91; 7.81; 15.63; 31.25; 62.50 and 125.00 µg/m. (Figure 2).

Cytotoxic effect of ethanol extract from *B. virgate* leaves was tested against T47D using colorimetric method MTT assay. All the cells were exposed to various concentration: 1.95; 3.91; 7.81; 15.63; 31.25; 62.50 and 125.00 µg/ml (Figure 3).




Figure 1. Cytotoxicity of ethanolic extracts of *B. virgate* leaves on the growth of HeLa cells were examined my MTT assay. Dose response curves constructed in the range 1.95 - 125.00 µg/ml after 24 h.




Figure 2. Cytotoxicity of ethanolic extracts of *B. virgate* leaves on the growth of WiDr cells were examined my MTT assay. Dose response curves constructed in the range 1.95 - 125.00 µg/ml after 24 h.




Figure 3. Cytotoxicity of ethanolic extracts of *B. virgate* leaves on the growth of T47D cells were examined my MTT assay. Dose response curves constructed in the range 1.95 - 125.00 µg/ml after 24 h.

Cytotoxic effect of ethanol extract from *B. virgate* leaves was tested against Vero using colorimetric method MTT assay. All the cells were exposed to various concentration: 1.95; 3.91; 7.81; 15.63; 31.25; 62.50 and 125.00 µg/ml (Figure 4).




Figure 4. Cytotoxicity of ethanolic extracts of *B. virgate* leaves on the growth of Vero cells were examined my MTT assay. Dose response curves constructed in the range 1.95 - 125.00 µg/ml after 24 h.

The IC<sub>50</sub> was determined based on concentration that induced 50% inhibition on the growth of the treated cells as compared to the untreated cells in triplicate after 24 h treated.

**Table 1. IC<sub>50</sub> and selectivity of ethanolic extract of *B. virgate* on various cell lines.**

Cancer cell line	IC <sub>50</sub> (ppm)	Selectivity Index (SI)
		(IC <sub>50</sub> of cancer cell line / IC <sub>50</sub> of Vero)
HeLa	18.991 ± 0.234	0.844
WiDr	18.925 ± 1.277	0.847
T47D	12.732 ± 0.945	1.258
Vero	16.022 ± 0.663	1.000

## DISCUSSION

The treatment of oral cancer relies on surgery, radiotherapy, chemotherapy or a combination of these methods.<sup>13</sup> Poor survival rates still occur, particularly for patients in advanced stages of the disease.<sup>14</sup> Natural products display a wide range of diversity in terms of their chemical structures and pharmacological properties. Several important antitumor drugs have been isolated from plants.<sup>15</sup>

The *B. virgata* of the genus *Boehmeria* have been widely studied by several author, in search of answers to their cytotoxic effect but no *B. virgata* was reported for each activity.

As the SI demonstrates the differential activity of extract, the greater the SI value is, the more selective it is. An SI value less than 2 indicates general toxicity of the pure compound<sup>16</sup>. Based on this, the SI data shown in Table 1 indicate that the extract was non selective in this cancer cell line, especially in HeLa, WiDr and T47D. This suggests its general toxicity to the cell.

Boehmeriasins A and B, new phenanthroquinolizidine alkaloids isolated from the aqueous ethanolic extract of *Boehmeria* genus, have cytotoxic activity against 12 cell lines.<sup>17</sup> Boehmeriasin A inhibits the proliferation of breast cancer cell MDA-MB-231 via the G1 phase cell cycle arrest and differentiation induction.

## CONCLUSIONS

In conclusion, we highlight the significant cytotoxic effect of *B. virgata* from the leaves, which introduces promising expectations for new projects in chemistry, pharmacology and toxicology although it was found to be non selective in HeLa, WiDr and T47D cell lines. These results may aid in achieving the development of an anticancer medicine obtained from the rain forest.

## REFERENCES

1. Ferlay, J., Shin, H.R., Bray, F., Forman, D., Mathers, C. and Parkin, D.M. 2010a. GLOBOCAN 2008 v1. 2, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10 [Internet].
2. Ferlay, J., Shin, H.R., Bray, F., Forman, D., Mathers, C. and Parkin D.M. 2010b. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Intl. J. Cancer*. **127**, 2893-28291
3. Newman, D.J., Cragg, G.M. and Snader, K.M. 2003. Natural products as sources of new drugs over the period 1981-2002. *J. Nat. Prod.* **66**, 1022-1037.
4. Ramawat, K.G. and Goyal, S. 2008. The Indian herbal drugs scenario in global perspectives, In Ramawat KG, Merillon JE "Bioactive Molecules and Medicinal Plants, Springer, Berlin Heidelberg, New York, 323.
5. Manggau, M., Hasan, H., Wahyudin, E. and Haryono, K. 2011. Efek Farmakologi Tanaman Antikanker yang Digunakan oleh Masyarakat Sulawesi Selatan. In: Mufidah and Lukman. Balitbangda Sulawesi Selatan, 2011, ISBN 978 602 8400 55 8.
6. Brands, S.J. 2007. *Systema Naturae 2000 The taxonomicon*. Universal Taxonomic Services, The Netherland. Cox, P.A. 1994. Ciba Foundation Symposium, Chichester, John Wiley & Sons. pp. 25-44.
7. Yan, J., Luo, D., Luo, Y., Gao, X., and Zhang, G. 2006. Induction of G1 arrest and differentiation in MDA-MB-231 breast cancer cell by boehmeriasin A, a novel compound from plant. *Intl. J. Gynecol Cancer*. **16**, 165-170.
8. Manggau, M., Mufidah and Lindequist, U. 2009. Antiproliferation Against Human Bladder Cancer 5637 Cell Line and Antioxidant Activity of Various Plant Extracts, *The Indonesian J. Nat. Prod.* **6**, 247-250
9. Mesia, G.K., Tona, G.L., Nanga, T.H., Cimanga, R.K., Apers, S., Cos, P., Maes, L., Pieters, L., and Vlietinck, A.J. 2008. *Antiprotozoal and cytotoxic screening of 45 plant extracts from Democratic Republic of Congo*, *J. Ethnopharmacol.* **115**(3), 409-415
10. Marshall, N.J., Goodwin, C.J. and Holt, S.J. 1995. A critical assessment of the use of microculture tetrazolium assays to measure cell growth and function. *Growth Regul* **5**, 69-84.
11. van Meerloo, J., Kaspers, G.J. and Cloos, J. 2011. Cell sensitivity assays: the MTT assay, *Methods Mol. Biol.* **731**, 237-245.
12. Shah, G.V., Muralidharan, A., Thomas, S., Gokulgandhi, M., Mudit, M., Khanfar, M. and El Sayed, K. 2009. Identification of a small molecule class to enhance cell-cell adhesion and attenuate prostate tumor growth and metastasis, *Mol. Cancer Ther.* **8**, 509-520.
13. Hsu, S., Singh, B., Schuster, G. 2004. Induction of apoptosis in oral cancer cells: agents and mechanisms for potential therapy and prevention. *Oral Oncol.* **40**, 461-473
14. Yao, M., Epstein, J.B., Modi, B.J., Pytynia, K.B., Mundt, A.J. and Feldman, L.E. 2007. Current surgical treatment of squamous cell carcinoma of the head and neck. *Oral Oncol.* **43**, 213-223.
15. Wall, M.E. and Wani, M.C. 1995. Camptothecin and taxol: discovery to clinic-thirteenth Bruce F. Cain Memorial Award Lecture. *Cancer Res.* **55**, 753-760.
16. Koch, A., Tamez, P., Pezzuto, J., and Soejarto D. 2005. Evaluation of plants used for antimalarial treatment by the Massai of Kenya. *J. Ethnopharmacol.* **101**, 95-99.
17. Luo, Y., Liu, Y., Luo, D., Gao, X., Li, B. and Zhang, G. 2003. Cytotoxic alkaloids from *Boehmeria siamensis*, *Planta Med.* **69**, 842-845.

