

Comparison of Shear Stresses of Different Fluid Flows Arising in the Boundary Layer Theory

Mohammad Riazuddin Molla

Department of Mathematics, Dhaka University, Dhaka-1000, Bangladesh

(Received: 28 July 2013; Accepted: 3 February 2014)

Abstract

The existence of positive solution for the singular second-order nonlinear boundary value problem arising in the boundary layer theory for the strong suction is studied. Then we compared the shear stress of the strong suction with the shear stress of the Homann flow. Also we compared the shear stress of the strong suction with the shear stress of the convergent flow.

Keywords : Suction, Positive solution, Homann flow, convergent flow.

I. Introduction

The differential equation

$$f''' + \alpha ff'' + \beta(1 - f'^2) = 0 \tag{1.1}$$

with boundary conditions

$$\left. \begin{aligned} f(\eta) = f'(\eta) = 0 \text{ at } \eta = 0 \\ f' \rightarrow 1 \text{ as } \eta \rightarrow \infty \end{aligned} \right\} \tag{1.2}$$

is known as Falkner-Skan boundary layer equation¹.

For $\alpha = 1$ and $\beta = 0.5$, the equation (1.1) takes the following form

$$f''' + ff'' + 0.5(1 - f'^2) = 0, \tag{1.3}$$

with boundary conditions (1.2). This equation represents Homann flow². For $\alpha = 0$ and $\beta = 1$, the equation (1.1) takes the following form

$$f''' - f'^2 + 1 = 0, \tag{1.4}$$

with boundary conditions (1.2). This equation represents flow in a convergent channel¹.

Shin² studied the differential equation (1.3) with boundary conditions (1.2) by using the constructive method such as the method of upper and lower solutions. In this article the constructive method such as the method of upper and lower solutions is also used to study the differential equation

$$\varphi''' + \varphi'' = 0 \tag{1.5}$$

with boundary conditions

$$\left. \begin{aligned} \varphi(\eta) = \varphi'(\eta) = 0 \text{ at } \eta = 0 \\ \varphi' \rightarrow 1 \text{ as } \eta \rightarrow \infty \end{aligned} \right\}, \tag{1.6}$$

which arises in the boundary layer theory for strong suction³.

Let the shear stress $g(x) = \varphi''(\eta)$ be the dependent variable and the tangential velocity $x = \varphi'(\eta)$ be the independent variable. The quantities x and g are called Crocco variables⁴.

Now
$$g' = \frac{dg}{dx} = \frac{d\eta}{dx} \frac{d}{d\eta} \left(\frac{d}{d\eta}(\varphi'') \right) = \frac{\varphi'''}{\varphi''}, \text{ which}$$

implies

$$g' \varphi'' = \varphi'''. \tag{1.7}$$

Differentiating (1.7) with respect to η one gets

$$\varphi^{iv} = g^2 g'' + g(g')^2.$$

As before differentiating (1.5) with respect to η one gets

$$\varphi^{iv} + \varphi''' = 0.$$

This gives

$$gg'' + (g')^2 + g' = 0.$$

Now
$$g(x) = \varphi''(\eta) = \frac{d}{d\eta}[\varphi'(\eta)].$$

Therefore

$$g(1) = \left[\frac{d}{d\eta}(\varphi'(\eta)) \right]_{x=1} = \left[\frac{d}{d\eta}(\varphi'(\eta)) \right]_{\varphi'(\eta)=1} = 0.$$

Again

$$g'(x) = \frac{dg}{dx} = \frac{\varphi'''}{\varphi''} = \frac{-\varphi''}{\varphi''}, \text{ which implies}$$

$$g'(0) = \frac{dg}{dx} \Big|_{x=0} = \frac{dg}{dx} \Big|_{\varphi'(\eta)=0} = \frac{dg}{dx} \Big|_{\eta=0} = \frac{-\varphi''(0)}{\varphi''(0)} = -1.$$

Thus by letting $g = \varphi''(\eta)$ and $x = \varphi'(\eta)$, equation (1.5) with boundary conditions (1.6) can be transformed into a second order singular nonlinear boundary value problem singularity at $x = 1$:

$$\left. \begin{aligned} gg'' + (g')^2 + g' = 0, \quad 0 < x < 1 \\ g'(0) = -1 \text{ and } g(1) = 0. \end{aligned} \right\} \tag{1.8}$$

For $\alpha = 1$ and $\beta = 0.5$, equation (1.1) with boundary conditions (1.2) can be written as

$$\left. \begin{aligned} g^2 g'' - \frac{1}{2}(1-x^2)g' = 0, \quad 0 < x < 1 \\ g'(0) = -0.5 \text{ and } g(1) = 0. \end{aligned} \right\} \tag{1.9}$$

by letting $g = f''(\eta)$ and $x = f'(\eta)$ and its positive solution has been studied by Shin². Shin² did not state the details of this transformation. In the present paper the details transformation of (1.8) from (1.5) with boundary conditions

*Author for correspondence. e-mail: mriaz_math@yahoo.com

(1.6) are shown using Crocco variables. The nonlinear differential equation (1.1) with boundary conditions (1.2) has been studied by many authors^{4,14} for different values of α and β using different methods.

For $\alpha = 0$ and $\beta = 1$, equation (1.1) with boundary conditions (1.2) can be written as

$$\left. \begin{aligned} g^2 g'' + (g')^2 - 2x &= 0, 0 < x < 1 \\ g'(0) = -0.86603 \text{ and } g(1) &= 0. \end{aligned} \right\} \quad (1.10)$$

by letting $g = f''(\eta)$ and $x = f'(\eta)$ and its positive solution has been studied by Molla⁹.

The equation (1.8) is equivalent to the nonlinear differential equation (1.5) with boundary conditions (1.6). So the positive solution of (1.8) on $[0, 1]$ is equivalent to the the shear stress $f''(\eta)$ on $[0, \infty)$.

The main purposes of this article are as follows:

- i) To establish the existence of positive solution of (1.8) using the constructive method such as the method of upper and lower solutions.
- ii) To compare the shear stress of the strong suction with the shear stress of the Homann flow.
- iii) To compare the shear stress of the strong suction with the shear stress of the convergent flow.

Definition 1.1. A function $\alpha_1 \in C^2[0,1]$ is called a positive upper solution of (1.8), if

$$\begin{aligned} \alpha_1 &> 0 \text{ on } (0,1) \\ \alpha_1 \alpha_1'' + (\alpha_1')^2 + \alpha_1' &\leq 0 \text{ on } (0,1) \\ \alpha_1'(0) &\leq -1 \text{ and } \alpha_1(1) \geq 0. \end{aligned}$$

Definition 1.2. A function $\alpha_2 \in C^2[0,1]$ is called a positive lower solution of (1.8), if

$$\begin{aligned} \alpha_2 &> 0 \text{ on } (0,1) \\ \alpha_2 \alpha_2'' + (\alpha_2')^2 + \alpha_2' &\geq 0 \text{ on } (0,1) \\ \alpha_2'(0) &\geq -1 \text{ and } \alpha_2(1) \leq 0. \end{aligned}$$

Similar definitions hold for positive upper and lower solutions of a perturbation (2.1) which will be given in the following section.

Definition 1.3. A function $g \in C[0,1] \cap C^2[0,1]$ is called a positive solution of (1.8), if

$$\begin{aligned} g &> 0 \text{ on } (0,1) \\ gg'' + (g')^2 + g' &= 0 \text{ on } (0, 1) \\ g'(0) &= -1 \text{ and } g(1) = 0. \end{aligned}$$

II. Existence of Positive Solution

Consider the nonlinear boundary value problem

$$\left. \begin{aligned} gg'' + (g')^2 + g' &= 0, 0 < x < 1 \\ g'(0) = -1 \text{ and } g(1) &= \frac{1}{p} \end{aligned} \right\} \quad (2.1)$$

for each $p \geq 1$, which may be viewed as a perturbation of (1.8).

To prove the existence of positive solution of (1.8) it is sufficient to established the existence of positive solution of (2.1).

Lemma 2.1. $g_{lp}(x) = 0.01x^\alpha \sqrt{1-x} + \frac{1}{p\alpha}$ is a positive

lower solution of (2.1), for each $p \geq 1$, where $\alpha = 10^k$ and k is integer, finite but very large.

Proof. It is clear that $g_{lp}(x) > 0$ on $(0, 1)$, $g'_{lp}(0) = 0$,

which can be written as $g'_{lp}(0) = 0 \geq -1$, $g_{lp}(1) = \frac{1}{p\alpha}$

, which can be written as $g_{lp}(1) = \frac{1}{\alpha p} \leq \frac{1}{p}$ and

$$h(x) = gg'' + (g')^2 + g' = g_{lp} g''_{lp} + (g'_{lp})^2 + g'_{lp} \geq 0,$$

for $0 < x < 1$ and $p \geq 1$.

Thus g_{lp} is a positive lower solution of (2.1). Consequently

$g_l = 0.01x^\alpha \sqrt{1-x}$ is a positive lower solution of (1.8). Notice that if $k=1$ then $h(x)$ is true for $(0, 0.9]$, if $k=2$ then $h(x)$ is true for $(0, 0.99]$, if $k= 3$ then $h(x)$ is true for $(0, 0.999]$ and so on.

Lemma 2.2. $g_{up}(x) = 2\sqrt{1-x} + \frac{1}{p}$ is a positive upper solution of (2.1) for each $p \geq 1$.

Proof. It is clear that $g_{up}(x) > 0$ on $(0, 1)$, $g'_{up}(0) = -1$, which can be written as $g'_{up}(0) = -1 \leq -1$,

$$g_{up}(1) = \frac{1}{p} \geq \frac{1}{p} \text{ and}$$

$$\begin{aligned} g_{up} g''_{up} + (g'_{up})^2 + g'_{up} \\ = -\{2\sqrt{1-x} + \frac{1}{p}\} \{0.5(1-x)^{-\frac{3}{2}}\} + \frac{1}{1-x} - \frac{1}{\sqrt{1-x}} \end{aligned}$$

≤ 0 , for $0 < x < 1$ and $p \geq 1$.

Thus g_{up} is a positive upper solution of (2.1).

Consequently $g_u = 2\sqrt{1-x}$ is a positive lower solution of (1.8).

Hence the following Lemma can be formulated from an application of Schauder's Fixed Point Theorem¹⁵.

Lemma 2.3. For any $p \geq 1$, there exists a positive solution $g_p \in C^2[0,1]$ of the problem (2.1) such that $g_{lp} \leq g_p \leq g_{up}$ on $0 \leq x \leq 1$, where g_{lp} and g_{up} are as given in Lemma 2.1 and Lemma 2.2 respectively.

Lemma 2.4. $g_p = (1-x) + \frac{1}{p}$ is a positive solution of (2.1)

for each $p \geq 1$.

Proof. It is clear that $g_p(x) > 0$ on $(0,1)$,

$$g'_p(0) = -1, g_p(1) = \frac{1}{p} \text{ and}$$

$$gg'' + (g')^2 + g' = g_p g''_p + (g'_p)^2 + g'_p = 0, \text{ for } 0 < x < 1 \text{ and } p \geq 1.$$

Thus g_p is a positive solution of (2.1). Consequently $g = 1-x$ is a positive solution of (1.8).

Thus by using Lemma 2.3 we can conclude that there exists a positive solution $g \in C^2[0,1]$ of the problem (1.8) such that $g_l \leq g \leq g_u$ on $0 \leq x \leq 1$, where g_l, g and g_u are as given above.

III. Results and Discussion

Here the positive solution of (1.8) is compared with the positive solution of (1.9) and (1.10) studied by Shin² and Molla⁹ respectively. The positive solution g_s of (1.9) lies between the positive lower and upper solutions $g_{sl} = 0.5(1-x)$ and $g_{su} = 2\sqrt{1-x}$ respectively obtained by Shin². The positive solution of (1.8) is

$g = 1-x$. The positive solution $g_c = \frac{\sqrt{4-6x+2x^3}}{3}$ of (1.10) lies between the positive lower and upper solutions

$$g_{lp} = \sqrt{1.3334 - 2.0000705x + 0.0000035x^2 + 0.6666667x^3}$$

and $g_{up} = 3\log(2-x) + 3$ respectively obtained by Molla⁹. Molla⁹ also showed that the positive solution converges to the positive lower solution.




Fig. 1. Different solutions such as (i) positive solution g obtained from present study (ii) positive lower solution g_l obtained from present study and (iii) positive upper solution g_u obtained from present study.




Fig. 2. Different solutions such as (i) positive solution g obtained from present study (ii) positive solution g_c obtained by Molla⁹ (iii) positive lower solution g_{sl} obtained by Shin² and (iv) positive upper solution g_{su} obtained by Shin².

Table 1. Numerical values of different positive, positive lower and positive upper solutions.

x	$g = 1 - x$	$g_c = \frac{\sqrt{4 - 6x + 2x^3}}{3}$	$g_{sl} = \frac{1}{2}(1 - x)$	$g_l = 0.01x^{\alpha} \sqrt{1-x}$ For $k=1$	$g_{su} = g_u = 2\sqrt{1-x}$
0	1	1.1547	0.50	0	2.000
0.2	0.8	0.9688	0.40	9.1589E-10	1.789
0.4	0.6	0.7589	0.30	8.1222E-10	1.549
0.6	0.4	0.5266	0.20	3.8242E-05	1.265
0.8	0.2	0.2733	0.10	4.8019E-04	0.894
1.0	0	0	0	0	0

From the above figure 1 and Table1 it is evident that the positive solution g of (1.5) obtained from present study lies on between g_l and g_u . Shin² did not find out the closed form positive solution of (1.9), only mention that there will be exist a positive solution of (1.9) lies on between g_{sl} and g_{su} . Since the positive solution converges to the positive lower solution so the numerical value of the positive solution g_s of (1.9) will be very close to the numerical value of g_{sl} . Hence from the above figure 2 it is apparent that the positive solution g of (1.5) is less than or equal to the positive solution g_c of (1.10) and the positive solution g_s of (1.9) less than or equal to the positive solutions g and g_c of (1.5) and (1.10) respectively. Notice that equality occurs only when $x=1$. From Table1 it is clear that the shear stress for the strong suction is decreasing uniformly with increasing tangential velocity. On the other hand the shear stress for the convergent flow is decreasing not uniformly with increasing tangential velocity.

IV. Conclusion

An outcome of the present study is to find a positive lower solution of equation (1.5). The existence of positive solution of (1.5) is established by using the constructive method such as the method of upper and lower solutions that is also an outcome. There exists a positive solution of (1.5) which lies on between the positive lower and upper solutions is shown. It is found that the shear stress for the strong suction is less than the shear stress for convergent flow with respect to the tangential velocity. It is also found that the shear stress for the Homann flow is less than the shear stress for convergent flow and strong suction with respect to the tangential velocity.

References

1. Shanti Swarup, 2000. Fluid Dynamics, Krishna Prakashan Media (P) Ltd. Merut, 622, 632,633.
2. Shin J.Y., 1997. A Singular nonlinear differential equation arising in the Homann flow, J. Math. Anal. Appl. **212**, 443-451.
3. Schlichting H., and K. Gersten,1999. Boundary Layer Theory, Springer, 113,114.
4. Soewono E., K. Vajravelu and R.N. Mohapatra , 1991. Existence and nonuniqueness of solutions of a singular nonlinear boundary-layer problem, J. Math. Anal. Appl. **159**,251-270.
5. Callegari A. and M. Friedman,1968. An analytical solution of a nonlinear, singular boundary value problem in the theory of viscous fluids, J. Math. Anal. Appl. **21**, 510-529.
6. Vajravelu K. ,E. Soewono and R.N. Mohapatra,1991.On solutions of some singular, non-linear differential equations arising in boundary layer theory, J. Math. Anal. Appl. **155**, 499-512.
7. Molla M.R., 2005. Existence and Uniqueness of positive solution of the suction of the fluid from the boundary layer, J. Math. and Math. Sci, JU, Savar, Bangladesh, **20**, 31-40.
8. Molla M.R.,2008. A singular non-linear BVP arising in the boundary layer flow along a flat plate, Ganit: J. Bangladesh Math. Soc. **28**, 59-67.
9. Molla M.R., 2012. A singular non-linear boundary value problem arising in a convergent channel, Bangladesh Journal of Physics, **12**, 15-26.
10. Molla M.R. and M. Begum, 2012. Existence and Uniqueness of positive solution of the injection of the fluid into the boundary layer.Jahangirnagar J.of Math.and Math.Sciences, **27**,103-114.
11. Molla M.R., 2013. An analytic treatment of the Falkner-Skan boundary layer equation. Journal of Science, University of Dhaka, **61(1)**, 139-144.
12. Molla M.R., M.K. Jaman and M. Hasan, 2011. Comparison of positive solutions for two boundary value problems arising in the boundary layer flow. Journal of Science, University of Dhaka, **59 (2)**, 167-172.
13. Molla M.R. and S.Banu, 2006. Existence and uniqueness of positive solution of a singular nonlinear BVP. Journal of Science, University of Dhaka, **.54 (2)**, 191-195.
14. Molla M.R. and S. Banu, 2003. Some singular nonlinear BVPS arising in the boundary layer flow. Ganit: Journal of Bangladesh Mathematical Soc., **23**, 91-103.
15. Schmidt K.,1970. A nonlinear boundary value problem, J. Differential Equations **7**, 527-537.