

Dissipation of Cypermethrin in Bean and Cauliflower

Nilufar Nahar^{1*}, Mohammad Motaher Hossain^{1,2}, Md. Nashir Uddin Al Mahmud¹, Mohammad Shoeb¹, Gulshan Ara Latifa² and Kamal Humayun Kabir³

¹Department of Chemistry, Dhaka University, Dhaka, Bangladesh

²Department of Zoology, Dhaka University, Dhaka, Bangladesh

³Entomology Division, Bangladesh Agricultural Research Institute, Gazipur, Bangladesh

(Received: 7 December 2015; Accepted: 24 January 2016)

I. Introduction

Cypermethrin [(RS)- α -cyano-3-phenoxybenzyl (1RS, 3RS; 1RS, 3RS)-3-(2, 2-dichlorovinyl) -2,2-dimethylcyclopropanecarboxylate] is one of the synthetic pyrethroids which have become most important insecticides in wide-scale use. Pyrethroids were reported to have relatively low toxicity as compared to organochlorine, organophosphorous, and carbamates pesticides¹ and their metabolites are also low toxic or non toxic compared with parent pyrethroid compounds². These are effective against the pests which are registrant to other group pesticides³ and account one-third of world insecticides use⁴ despite their high cost.

The vegetables are low in fat, high in dietary fibers; contain minerals and vitamins, possessing a very high nutritional density. But repeated applications and over doses of pesticides in the vegetable cultivation by the farmers make these unsafe and health hazardous for every day consumption. It is necessary to monitor and assess of the pesticide contamination in vegetables samples. In continuation of our work on vegetables, we are now reporting dissipation patterns of cypermethrin in cauliflower (*Brassica oleracea* var.) and bean (*Phaseolus vulgaris*) and their safe harvesting periods.

II. Experimental

The experiments was conducted at the experimental field of Bangladesh Agricultural Research Institute, Joydebpur, Gazipur situated at latitude 23° 46' N and longitude 90° 23' E with an elevation of 8.45 meter the sea level. Seeds were sown in the nursery bed at research field of Entomology Division, Bangladesh Agricultural Research Institute, Gazipur. The plants were lightly irrigated regularly for ensuring proper growth and development of the seedlings. Forty days old seedlings were transplanted in the well prepared experimental plot in 28th October 2012. "Ripcord-10-EC" (100 g cypermethrin per liter) was sprayed at recommended dose of (1 mL in 1 L water = 100 mg cypermethrin in 1 kg water) in experimental bed (cauliflower at 2 December & bean at 18 December, 2012). Before spraying, blank/control samples were collected. Then two replicate samples from each bed were collected at the intervals of 0 (after 2 hours of spraying), 1, 3, 5, 7, 10 and 15 days after spraying. They were kept in a freezer by wrapping with clean polythene bag (jeep-lock) at temperature below -20 °C.

Edible part of each vegetable sample (250 g) was cut into small pieces and homogenized by means of a kitchen blender. 20 mL ethyl acetate was added to 10 g homogenized sample in 50 mL Teflon tube and shaken for 1 minute by hand & vortexed for 1 min. Anhydrous MgSO₄ (6 g) & 1.5 g NaCl were added and vortexed for 1 min and then centrifuged for 5 minutes at 4000 rpm. 10 mL supernatant was taken in 100 mL round bottomed flask, evaporated in rotary evaporator and then reconstituted in n-hexane (2 mL). A glass column (40 cm long and 12 mm internal diameter) was packed with a 10.5 g mixture of aluminum oxide, florisil and charcoal (10:10:1) in n-hexane. The column was equilibrated with 50 mL n-hexane and then the sample extract in n-hexane (2 mL) was applied to the column. The column was washed with 20 mL of n-hexane and eluted with 100 mL of dichloromethane at the rate of 1 mL min⁻¹. The eluent was concentrated to dryness by a rotary vacuum evaporator and re-dissolved the residue in 2 mL of n-hexane and analyzed by GC-ECD.

A Shimadzu 17A GC system equipped with Electron Capture Detector was used for analysis. The injector & detector temperatures were 280 °C & 300 °C. Separations were performed on HP-5MS quartz capillary column (30 m × 250 μ m *i.d.* and 0.25 μ m film thickness), nitrogen was used as a carrier gas and flow rate was 1.0 mL min⁻¹. Samples were injected manually in split-less split mode and injection volume was 1 μ L. The oven temperature was programmed as initial temperature of 120 °C held for 2 min, increased at 10 °C min⁻¹ to 270 °C; held for 1 min. and then another increased at 2 °C min⁻¹ to 290 °C; held for 3 min.

III. Results and Discussion

The residual level of cypermethrin in bean and cauliflower was analyzed by GC-ECD using external standard calibration method. The linear calibration curves over six calibration levels, from 0.025-2.0 mg L⁻¹ were constructed by the direct injection of calibration standards. The linearities were excellent with a correlation coefficient of $r^2 = 0.998$. The unnecessary compounds interfering with the analytes were examined by comparing the chromatograms of the standard, blank sample and spiked sample. There were no interference peaks at the retention time of cypermethrin. The LOD (limit of detection; Signal to noise ration, 3:1) and

*Author for correspondence. e-mail: nilufarnahar@yahoo.com

LOQ (limit of quantification; Signal to noise ratio, 10:1) were found to be 0.01 & 0.033 mg kg⁻¹, respectively. The recovery experiments were performed in three replicates at 2 spiking levels (0.25 & 1.0 mg kg⁻¹). The spiked samples were left to stand for 2 h to allow for the adsorption of pesticide onto the samples. The recoveries were 76-107% with RSD ≤ 13% which were in accordance with Codex Alimentarius⁵. Residual amounts of cypermethrin in all the analyzed samples in which recommended doses sprayed are presented in **Table 1**. The pesticide was degraded with time following first order kinetics (**Fig. 1**). The residue

concentration of cypermethrin in bean was found to be 3.8 ± 0.53 and 0.32 ± 0.01 mg kg⁻¹ at 0 day (2 hours after spray) and at 15 days after application, respectively. Residue in cauliflower was found to be 6.7 ± 1.8 at 0 and 0.12 ± 0.01 mg kg⁻¹ at 15 days, respectively after application. Residue of cypermethrin in bean and cauliflower was found to be below MRL on 10 and 7 days after application, respectively. The results are in agreement with the findings of Chai *et al* (2009)⁶ and Nahar *et al.* (2012)⁷ where dissipation of cypermethrin was studied in green mustard and tomato, respectively.

Table 1. Cypermethrin residues in vegetables matrix at different time intervals following its application			
Sample	Day after spraying	Bean [(Av. ± SD), mg kg ⁻¹]	Cauliflower [(Av. ± SD), mg kg ⁻¹]
Control	-	Not detected	Not detected
Day	0	3.77 ± 0.53	6.68 ± 1.86
	1	2.69 ± 0.19	2.94 ± 0.16
	3	1.13 ± 0.08	1.72 ± 0.05
	5	0.94 ± 0.11	1.12 ± 0.04
	7	0.89 ± 0.06	0.67 ± 0.05
	10	0.45 ± 0.11	0.17 ± 0.03
	15	0.32 ± 0.01	0.13 ± 0.01




Fig 1. Dissipation curves of cypermethrin in (a) bean and (b) cauliflower

IV. Conclusion

Farmers use cypermethrin for bean and cauliflower very frequently, even twice a day and harvest crop immediately after pesticide application. Our finding suggests that farmers should apply cypermethrin in bean and cauliflower as recommended dose and harvest after 7-10 days for consumer safety.

Acknowledgment

The authors are grateful to International Science Programme (ISP), Uppsala, Sweden for financial supports.

References

1. Columé A, S Cardenas, M Gallego, M Valcarcel, 2001. Semiautomatic multiresidue gas chromatographic method for the screening of vegetables for 25 organochlorine & pyrethroid pesticides. *Analytica Chimica Acta*, **436** (1), 153-162.
2. Chen ZM and YH Wang, 1996. Chromatographic methods for the determination of pyrethrin and pyrethroid pesticide residues in crops, foods and environmental samples. *Journal of Chromatography A*, **754**(1-2), 367-395.
3. The Pesticide Manual, Fourteenth Edition 2006. Editor: C D S Tomlin, Publisher: BCPC (British Crop Production Council).
4. Fundamental concepts of Environmental Chemistry, G.S. Sodhi, Department of Chemistry, S.G.T.B. Khalsa College, University of Delhi, 2002.
5. Codex Alimentarius, Guidelines on good laboratory practice in pesticide residue analysis CAC/GL40-1993.
6. Chai L. K, N. Mohd-Tahir, H. C. B. Hansen, 2009. Year *Pest Management Science*, **65** (2), 189-196.
7. Nahar N, M Shoeb, M I R Mamun, S Ahmed, MM Hasan and A Kabir, 2012. Studies of Dissipation Pattern of Cypermethrin in Tomato. *Journal of Bangladesh Chemical Society*, **25**(2), 200-203.

