

## ***tert.*-Butylation of *p*-Cresol with *tert.*-Butyl Chloride**

Md. Shahrzaman, Manoranjan Saha\*, Nusrat Sharmin, Md. Kabir Hossain, Md. Tafsir Uddin Bhuyan, A. M. Sarwaruddin Chowdhury, Dipti Saha, Md. Zahangir Alam and Mohammad Ismail  
Department of Applied Chemistry and Chemical Engineering, Dhaka University, Dhaka-1000, Bangladesh

Received on 25.02.2010, Accepted for Publication on 30.06.2010

### **Abstract**

2-*tert.*-Butyl-4-methylphenol was obtained in high yield by the alkylation of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride. The effects of the variation of molar ratio of *p*-cresol to *tert.*-butyl chloride, temperature, amount of AlCl<sub>3</sub> and time of reaction have been studied on the reaction to find out an optimum condition.

### **I. Introduction**

The outstanding antioxidants and multifunctional stabilizers for fuels, lubricating oils and polymeric materials are based on alkylphenols and cresols<sup>1-3</sup>. Alkyl cresols and some of their derivatives can also be used as herbicides, bactericides, insecticides etc.<sup>3,4</sup>. Alkylated cresols with an alkyl group of 9-12 carbon atoms are valuable intermediates for surfactants and nonionic detergents<sup>2,3,5</sup>.

Alkylated cresols have been obtained by several authors by alkylation of isomeric cresols with olefins<sup>6-15</sup> and alcohols<sup>16-25</sup> using different catalysts. But no attempt has so far been made to investigate the reactions of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride.

In the present work, attempt has been made to investigate the reaction of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride as catalyst.

### **II. Experimental**

The reactions were carried out in a three necked round bottomed flask fitted with a condenser, a thermometer, a dropping funnel and a magnetic stirrer. *p*-Cresol-AlCl<sub>3</sub> mixture was heated to the desired temperature and *tert.*-butyl chloride was introduced into the mixture gradually for a certain period of time (time of addition) with constant stirring. The reaction mixture was stirred for an extended period of time (time of stirring) at the same temperature after the complete addition of total amount of *tert.*-butyl chloride. The reaction mass was then cooled to room temperature, dissolved in petroleum ether, neutralized with saturated NaHCO<sub>3</sub> solution followed by washing with distilled water and subjected to distillation. Unreacted reactants and solvent were distilled off at atmospheric pressure. The residual product was finally distilled and characterized by spectral (IR, <sup>1</sup>H NMR) means.

### **III. Results and Discussion**

The results of the reaction of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride over the temperature range of 80 to 160°C are presented in Tables 1-4 and in Figs. 1-4. Molar ratio of *p*-cresol to *tert.*-butyl chloride was varied from 4:1 to 8:1, amount of AlCl<sub>3</sub> from 5 to 8% by wt. of *p*-cresol and reaction time from 1 to 4.5h. The *tert.*-butyl group substituted the aromatic ring to the *ortho*- position with respect to the -OH group and 2-*tert.*-butyl-4-methylphenol was obtained in good yield.

#### **The effect of variation of molar ratio of *p*-cresol to *tert.*-Butyl chloride**

The effect of variation of molar ratio of *p*-cresol to *tert.*-butyl chloride on the reaction of *p*-cresol with *tert.*-butyl chloride at temperature of 140°C, addition time of 2h and stirring time of 1h and in the presence of anhydrous aluminium chloride (5% by wt. of *p*-cresol) was studied when 62.1, 74.1 and 81.5% yield of 2-*tert.*-butyl-4-methylphenol were obtained at molar ratio of 4:1, 5:1 and 8:1, respectively. This showed that the yield of the product increased with the increase in molar ratio of *p*-cresol to *tert.*-butyl chloride. Thus the yield increased from 62.1 to 81.5% when the molar ratio of *p*-cresol to *tert.*-butyl chloride increased from 4:1 to 8:1, and it was maximum at a molar ratio of 8:1 (Fig. 1).

#### **The effect of variation of temperature**

The effect of variation of temperature on the reaction of *p*-cresol with *tert.*-butyl chloride in molar ratio of 5:1, addition time of 2h, stirring time of 1h and in presence of anhydrous aluminium chloride (6% by wt. of *p*-cresol) was studied when 62.5, 74.6, 80.3 and 83.6% yield of 2-*tert.*-butyl-4-methylphenol were obtained at temperature of 80, 100, 140 and 160°C, respectively. This indicated that the yield of the product (2-*tert.*-butyl-4-methylphenol) increased with the increase in temperature. The yield increased from 62.5 to 83.6% for the increase in temperature from 80 to 160°C and it was found maximum at 160°C (Fig. 2).

#### **The effect of variation of the amount of AlCl<sub>3</sub>**

The effect of variation of amount of anhydrous aluminium chloride on the reaction of *p*-cresol with *tert.*-butyl chloride at temperature of 140°C, molar ratio of 5:1, addition time of 2h, stirring time of 1h was carried out when 74.1, 80.3 and 84.9% yield of 2-*tert.*-butyl-4-methylphenol were obtained using the catalyst in an amount of 5, 6 and 8% by wt. of *p*-cresol, respectively. Thus the yield of the product was found to increase with the increase in the amount of AlCl<sub>3</sub>. The yield increased from 74.1 to 84.9% when the amount of anhydrous aluminium chloride was increased from 5 to 8%

\*Corresponding Author: E-mail: manoranjansaha2005@yahoo.com  
Tel: +880-2-8621770, Fax: +880-2-8615583

and it was observed to be maximum at 8% by wt. of *p*-cresol (Fig. 3).

### The effect of variation of reaction time

The effect of variation of reaction time on the reaction of *p*-cresol with *tert.*-butyl chloride (molar ratio of *p*-cresol to *tert.*-butyl chloride = 5:1) at temperature of 140°C in the presence of anhydrous aluminium chloride (6% by wt. of *p*-cresol) was investigated by three sets of experiments with different addition time and stirring time (Fig. 4). From the first set of experiments, it was observed that the yield of the product increased from 71.2 to 78.6% when addition time was varied from 1 to 2.5 h (Fig. 4, Curve 1). Second set of experiments showed that the best yield was obtained when addition time was 2.5h (Fig. 4, Curve 2). Finally the third set of experiments showed that the yield increased with the increase in stirring time (Fig. 4, Curve 3).

### Optimum conditions

Thus the following conditions were considered as optimum for the production of 2-*tert.*-butyl-4-methylphenol:

temperature = 140°C, molar ratio of *p*-cresol to *tert.*-butyl chloride = 5:1, amount of anhydrous aluminium chloride = 6% by wt. of *p*-cresol, addition time = 2 h and stirring time = 2.5 h.

### Spectral studies (IR, <sup>1</sup>H NMR) of 2-*tert.*-butyl-4-methylphenol


The IR spectrum of 2-*tert.*-butyl-4-methylphenol showed bands at 815.89 cm<sup>-1</sup> and 877.61 cm<sup>-1</sup> indicating the presence of 1, 2, 4 – trisubstituted benzene ring. The spectrum also showed absorption bands at 3387.00 cm<sup>-1</sup>, 3018.60 cm<sup>-1</sup>, 2924.09 cm<sup>-1</sup> and 1606.70 cm<sup>-1</sup> for –OH group, aromatic =C–H, aliphatic C–H and aromatic ring C–C stretching, respectively.

Table 1 shows the positions of signals of the protons in the <sup>1</sup>H NMR spectrum of 2-*tert.*-butyl-4-methylphenol.


**Table 1. Signals of the protons in the <sup>1</sup>H NMR spectrum of 2-*tert.*-butyl-4-methylphenol**

Observed signals of the protons	Chemical shift in δ ppm
Aromatic ring protons	6.55 - 7.24
One proton on the –OH group	4.76 – 5.01
Three protons of –CH <sub>3</sub> group	2.06– 2.42
<i>tert.</i> -Butyl group protons	1.24 – 1.80


2-*tert.*-Butyl-4-methylphenol had b.p. = 237 °C,  $n_D^{20} = 1.5398$  and  $d_4^{20} = 0.963$ .




**Fig. 1.** The effect of variation of molar ratio of *p*-cresol to *tert.*-butyl chloride on the reaction of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride (temperature = 140 °C, amount of anhydrous aluminium chloride = 5% by wt. of *p*-cresol, addition time = 2h and stirring time = 1h).



**Fig. 2.** The effect of variation of temperature on the reaction of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride (molar ratio of *p*-cresol to *tert.*-butyl chloride = 5: 1, amount of anhydrous aluminium chloride = 6% by wt. of *p*-cresol, addition time = 2h and stirring time = 1h).



**Fig. 3.** The effect of variation of the amount of AlCl<sub>3</sub> on the reaction of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride (temperature = 140°C, molar ratio of *p*-cresol to *tert.*-butyl chloride = 5:1, addition time = 2h and stirring time = 1h).



**Fig. 4.** The effect of variation of reaction time on the reaction of *p*-cresol with *tert.*-butyl chloride in the presence of anhydrous aluminium chloride (temperature = 140°C, molar ratio of *p*-cresol to *tert.*-butyl chloride =5:1 and amount of AlCl<sub>3</sub> = 6% by wt. of *p*-cresol):

Curve 1 -The effect of variation of addition time (stirring time = 0 h);

Curve 2 -The effect of variation of addition time (reaction time = 3 h);

Curve 3 -The effect of variation of stirring time (addition time = 2 h).

1. Ravikovich, A. M., 1964. Antioxidants for minerals and synthetic lubricating oils. *Chemistry and Technology for Fuels and Oils*, **11**, 68-71.
2. Shreve, R. N. and J. A. Brink, 1977. Chemical process industries. 4<sup>th</sup> Ed., (McGraw-Hill International Book Company, London). pp. 814.
3. Lebedev, N. N., 1984. Chemistry and Technology of Basic Organic and Petrochemical Synthesis, Mir Publishers, Moscow, Vol. 1 and 2, pp 638.
4. Melnikov, N. N., Y. A. Baskahov and K. S. Bokrev, 1954. Chemistry of herbicides and plant growth regulators. *Gkhi. Moscow*. **38**.
5. Dimitriev, S. A., K. D. Korener and O. N. Tsvetkov, 1961. Synthesis of detergents of *o*, *p* - type based on phenols derived from peat oils. *Torfyannaya. Prom.* **32 (6)**, 24-27.
6. Topchiev, A. V., Zavgorodnii S. V. & V. G. Kryuchkova, 1962. Alkylation of Organic Compounds with Olefins (Academy of Sciences USSR, Moscow). pp. 324.
7. Kharchenko, L. S., S. V. Zavgorodnii, 1964. Alkylation of *p*-cresol with propylene in the presence of BF<sub>3</sub>.H<sub>3</sub>PO<sub>4</sub> and BF<sub>3</sub>.Et<sub>2</sub>O. *U Kr. Khim. Zh.* **30 (2)**, 187-190.
8. Babakhianov, R. A., D. E. Mishev, Gaidorova and T. Samedova, 1965. Alkylation of cresols by olefins. *Azer. khim. Zh.* **1**, 42-44.

9. Gusev, V. K., V. I. Isagulayants, P. S. Belov and Ya. N. Shekhter, 1969. Continuous alkylation of *p*-cresol by isobutylene on Ku-2 ion exchange resin. *Khim. Prom.* **42**, 96-98.
10. Saha, M., S. K. Ghosh, S. K. Saha and M. A. B. Sarker, 1994. Alkylation of *p*-cresol with cyclohexene in the presence of sulphuric acid. *Bangladesh J. Sci. Ind. Res.* **29 (1)**, 144-148.
11. Saha, M., S. Saha, and M. Mosiuzzaman, 1996. Alkylation of *p*-cresol with cyclohexene in the presence of benzenesulfonic acid. *Indian journal of chemical Technology.* **3**, 292-294.
12. Saha, M., S. Chowdhury, M. A. B. Sarker, Y. N. Jolly and D. Saha, 1997. A study of alkylation of cresols with cyclooctene. *Bangladesh J. Sci. Ind. Res.* **32 (1)**, 63-66.
13. Saha, M., H. M. N. E. Mahmud, S. K. Ghosh, M. B. Zaman, D. Saha and S. K. Saha, 2000. Cycloalkylation of *p*-cresol with cycloalkenes in the presence of borontrifluoride etherate. *Bangladesh J. Sci. Res.* **18 (2)**, 273-276.
14. Saha, S. K., D. Saha and M. Saha, 2003. Cycloalkylation of *p*-cresol with cyclooctene. *Dhaka Univ. J. Sci.* **51 (2)**, 299-300.
15. Saha, M., H. C. Dey, M. Z. Karim, M. Ismail and D. Saha, 2008. Alkylation of cresols with cyclohexene in the presence of *p*-toluenesulphonic acid. *Bangladesh J. Sci. Ind. Res.* **43 (2)**, 277-282.
16. Abdurasuleva, A. R., N. Ismailov and Y. Yotdashev, 1969. Alkylation of cresols by cyclohexanol in the presence of KU-2 cation exchanger. *Zh. Obsh. Khim.* **13 (5)**, 50-52.
17. Saha, M. and R. K. Roy, 1988. Reaction of *p*-cresol with *n*-pentanol in the presence of sulphuric acid. *Bangladesh J. Sci. Ind. Res.* **6**, 83-90.
18. Saha, M. and R. K. Roy, 1989. Study of alkylation of *p*-cresol with *n*-hexanol. *Bangladesh J. Sci. Ind. Res.* **24 (14)**, 33-40.
19. Saha, M., Y. N. Jolly, D. Saha and M. A. B. Sarker, 1997. Alkylation of *p*-cresol with *n*-alcohol. *Bangladesh J. Sci. Ind. Res.* **32 (1)**, 67-69.
20. Saha, D. and M. Saha, 2002. Alkylation of *p*-xylene and *p*-cresol with cycloalcohol. *Bangladesh J. Sci. Ind. Res.* **37 (1-4)**, 113-122.
21. Rana, A.A., M. Kamruzzaman, M. A. B. Sarker, A. Alam and M. Saha, 2003. Benzylolation of cresols with benzyl alcohol in the presence of sulphuric acid. *Dhaka Univ. J. Sci.* **51 (2)**, 169-178.
22. Basak, P. K., S. K. Saha, M. Asaduzzaman and M. Saha, 2004. Alkylation of cresols with *tert.*-amylalcohol. *Dhaka Univ. J. Sci.* **52 (3)**, 287-292.
23. Saha, M., S. Biswas, S. K. Saha and R. F. Rafique, 2004. A mathematical model for the alkylation of *p*-cresol with *tert.*-amylalcohol in the presence of sulphuric acid. *Bangladesh J. Sci. Ind. Res.* **39 (3-4)**, 139-146.
24. Saha, M., S. Biswas, S. K. Saha and R. F. Rafique, 2006. Reaction of cresols with cycloalcohols. *Dhaka Univ. J. Sci.* **54 (1)**, 35-38.
25. Ismail, M., M. S. Jamal, S. T. A. Islam, M. Z. Alam, M. Ashaduzzaman and M. Saha, 2007. A mathematical model for the benzylolation of *p*-cresol with benzyl alcohol. *Bangladesh J. Sci. Ind. Res.* **42 (2)**, 187-194.

