

Original Article

Performance of Arbuscular Mycorrhiza Inoculated *Acacia mangium* Seedlings on Degraded Land with Different Rates of Phosphorus

M Abdus Satter^{1*}, M Musa Hanafi², Tenku MM Mahmud² and Hashim Azizah²

¹Soil Science Division, Bangladesh Agricultural Research Institute (BARI), Joydebpur, Gazipur 1701, Bangladesh, ²Faculty of Agriculture, University Putra Malaysia (UPM), Serdang, Selangor, Malaysia

[Received 08 May 2006; Accepted 21 April 2007]

The experiment was conducted in the degraded ex-tin mining farm of Universiti Putra Malaysia during November 1998 to May 1999. The experiment was laid out in split plot design with three replications. Three phosphorus (P) rates, viz., 0, 50 and 100 g Gafsa phosphate rock (GPR) per plant in combination with and without arbuscular mycorrhiza (AM) inoculation were studied on newly planted tree species *Acacia mangium*. Mycorrhiza inoculation was assigned to the main plot and P dose to the subplot. Three-month-old AM pre-inoculated seedlings were planted in the field at 2 m x 2 m spacing. The experiment was continued until 160 days after planting (DAP). Mycorrhiza inoculation and rate of P application significantly influenced growth and development of the *A. mangium* plants. The highest growth of un-inoculated *A. mangium* plants was observed with 100 g GPR per plant; whereas for AM inoculated plants the highest growth was observed with 50 g GPR per plant. This indicates that P requirement of newly planted *A. mangium* plants on degraded ex-tin mining land could be reduced to 50 g GPR per plant, i.e., by 50% with AM inoculation.

Keywords: Arbuscular mycorrhiza, *Acacia mangium*, Degraded land, Phosphorus

Introduction

Degraded ex-tin mining land covers vast areas in the tropics. Highly weathered acid soils in the tropics including ex-tin mining soils are widely deficient in phosphorus (P). Phosphorus applied to these soils is rapidly transformed into low solubility Al- or Fe-bound P. Up to 80% of the applied P might be lost because of transformation into such insoluble forms¹. Due to inefficient utilization of soluble superphosphates by plants in these soils, less soluble and less expensive phosphate rocks (PRs) have been suggested for crop production². Uptake of P from PRs might be improved through combined application with AM³.

The AM fungi are beneficial fungi forming symbiotic association with roots of the most plant species. The external AM hyphae extend several centimetres from the infected root surface and help in exploration of greater soil volume to absorb more nutrients and moisture from the soil. Besides, they increase the rate of photosynthesis of the host plants. They also enhance production of growth regulating substances in the host plants⁴. They improve P uptake from less soluble sources like phosphate rock (PR). They also help legumes in improving biological N₂-fixation through improved P nutrition⁵. Furthermore, they help plants in uptake of other nutrients as well⁶. Available reports suggest that low P level increases AM colonization and hence

growth and nutrition of plants. On the other hand, high P levels decrease root colonization, with plants having little or even no benefits out of mycorrhizal inoculation⁷. There are also evidences that under high soil P conditions AM can become parasitic and hence depress yield⁸. Soil P level and rate of P application are therefore crucial in influencing AM association and deriving benefits there of. Gafsa phosphate rock (GPR) in combination with AM was found to produce almost equivalent biomass of *A. mangium* seedlings to those with combined application of TSP and AM on the degraded ex-tin mining soils⁹. Such information suggest further study on determining optimum P dose (from GPR) for *A. mangium* seedlings grown on degraded ex-tin mining land. The present study was therefore undertaken to observe the effect of different P rates (from GPR) with and without AM inoculation on the growth performance of *A. mangium* in the degraded ex-tin mining land.

Materials and Methods

The experiment was conducted in the degraded ex-tin mining farm of Universiti Putra Malaysia during November 1998 to May 1999. The soil was deficient in phosphorus, potassium and magnesium; and calcium content was at the critical level. Organic matter content and soil pH was low. The physical and chemical characteristics of the soil are presented in Table 1.

*Corresponding author:

Dr. M Abdus Satter, Principal Scientific Officer (Soil), Bangladesh Agricultural Research Council (BARC), Farmgate, Dhaka 1215, Bangladesh
Tel (Office): (02) 9132411; Cell: 01716 420890; E-mail: satterbari@yahoo.com

Table 1. Initial physical and chemical properties of soils of the experimental field

Soil variable	Content	Soil variable	Content	Critical level
Sand (%)	79.30	K (meq/100g)	0.12	0.2
Silt (%)	14.60	Ca (meq/100g)	2.35	2.0
Clay (%)	6.10	Mg (meq/100g)	0.18	0.8
pH (H ₂ O)	5.56	CEC (meq/100g)	2.37	-
Organic C (%)	0.75	Zn (ppm)	13.11	2.0
N (%)	0.09	Cu (ppm)	38.27	1.0
P (ppm)(Bray II)	7.36	Mn (ppm)	17.27	5.0

The experiment was laid out in split plot design with three replications. Three P rates, *viz.*, 0, 50 and 100 g GPR per plant in combination with AM and without AM were studied. Gafsa phosphate rock (GPR) containing 13% P was used as the source of P. Mycorrhiza inoculation was assigned to the main plot and P dose to the subplot. Seedlings of *A. mangium* were raised in black polythene bags containing 1 kg fumigated tin-tailing sand. For AM treatment, the seedlings were inoculated with 50 g of mixed AM inoculum per bag per seedling. The seeds were inoculated with *Rhizobium* irrespective of AM and P treatments. Seedlings were transplanted to the field at the age of three months. The unit plots measured 8 m x 10 m. Pits of 0.5 m x 0.5 m x 0.5 m size were dug at 2 m x 2 m spacing. Fertilizers were applied into the pits. Phosphorus fertilizer was applied as per treatment. The other fertilizers were applied as blanket dose at the rate of 60, 20, 40 and 1 g per pit of MP, gypsum, dolomite and manganese chloride, respectively. Fertilizers were mixed well with the soils of the pits before refilling.

Seedlings were planted during the fourth week of November 1998, seven days after pit preparation. No additional AM inoculum was applied during planting seedlings. A life-saving irrigation was given at three days after planting. The land was covered with polythene mulch for controlling weeds. Side shoots were plucked weekly up to 2 m plant height. The experiment was continued until 160 days after planting (DAP). Data on plant

height and basal diameter (at 10 cm above ground level) were measured from all 20 plants in each plot at 10, 40, 70, 100, 130 and 160 DAP. Spad values were measured on the five randomly selected youngest fully expanded leaves from five plants per plot. At 150 DAP, root samples were collected from five randomly selected plants per plot. Rhizosphere soil samples of the same plants were also collected for counting AM spores. Five sub-samples around each plant were collected, bulked and mixed thoroughly to make a composite sample. To assess AM infection, roots were processed after Koske and Gemma¹⁰ and observed under compound microscope. A total of 100 readings were taken per sample and the infection was expressed in percentage. Spore numbers in rhizosphere soil were determined by wet sieving and decanting method¹¹. Data were analyzed using statistical package MSTAT-C¹².

Results and Discussion

Plant growth

Height and basal diameter growth of *A. mangium* on degraded ex-tin mining land was significantly influenced by AM inoculation and rate of P application. Interaction effect of AM inoculation and age of plant was significant both on plant height and basal diameter growth of *A. mangium* (Figure 1). Height and basal diameter growth of AM inoculated and un-inoculated plants were identical until 70 DAP but became significantly different from 100 DAP onwards. Interaction effect of AM and P dose on the plant height and basal diameter growth of *A. mangium* was also significant. The AM inoculated plants given 50 g GPR per plant produced significantly higher plant height (1.23 m) and basal diameter (1.47 cm) compared to those with other treatments (Table 2). The AM inoculated plants given 0, 50 and 100 g GPR per plant produced 1.12, 1.26 and 1.05 times higher plant height and 1.16, 1.27 and 1.08 times higher basal diameter, respectively over those of un-inoculated plants. Interaction effect of AM inoculation, rate of P application and plant age on the height and basal diameter growth of *A. mangium* was not found significant.




Figure 1. Interaction effect of arbuscular mycorrhiza (AM) and plant age on (a) plant height and (b) basal diameter of *Acacia mangium* on degraded ex-tin mining land. DAP = Days after planting

Table 2. Interaction effect of arbuscular mycorrhiza (AM) and phosphorus (P) dose on the height and basal diameter of *Acacia mangium* on degraded ex-tin mining land


P dose (g GPR per plant)	Plant height (m)		Basal diameter (cm)	
	With AM	Without AM	With AM	Without AM
0	1.02 ^{bcd}	0.91 ^d	1.21 ^{cd}	1.04 ^e
50	1.23 ^a	0.97 ^{cd}	1.47 ^a	1.16 ^d
100	1.12 ^{ab}	1.06 ^{bc}	1.38 ^b	1.27 ^c
F-test	*		**	

In a column, the figures having common letter/s do not differ significantly at 1 and 5% level of probability. GPR = Gapspha phosphate rock

Height and basal diameter growth indicates that 100 g GPR per plant may not be sufficient for un-inoculated *A. mangium* plants. But for the inoculated plants, 100 g GPR per plant might probably stretched beyond the threshold level for optimum mycorrhizal activity. Because high P level is reported to be detrimental to AM activity and hence growth of plants⁷. Several evidences on the growth enhancement of different tree species with AM inoculation under field conditions are also available¹³⁻¹⁴.

Spad value

Effect of AM inoculation and rate of P application on spad value of *A. mangium* leaves has been presented in Figure 2. Rate of P application and AM inoculation also significantly influenced spad value of the leaves of *A. mangium* grown on degraded ex-tin mining land. Effect of AM inoculation on spad value of *A. mangium* leaves was found significant. Spad value of AM inoculated plants (45) was significantly higher than that of un-inoculated plants (41). Rate of P application also had significant ($p \leq 0.05$) effect on spad value of *A. mangium* leaves. *Acacia mangium* plants given 50 and 100 g GPR per plant produced same spad value (50), which was significantly higher than that of control plants (40). Interaction effect of AM inoculation and rate of P application on spad value of *Acacia mangium* leaves was not found significant.

**Figure 2.** Effect of arbuscular mycorrhiza (AM) inoculation and phosphorus (P) rate on the spad value of *Acacia mangium* leaves.

Spad value is known to be an indirect measurement of chlorophyll content which indicates the photosynthetic efficiency of a leaf¹⁵. The inoculated plants in the present study had higher spad value

enhancing higher photosynthesis and thereby higher biomass production. It was observed in a previous study that *A. mangium* leaves with higher spad values had higher photosynthetic efficiency⁹. A positive relationship between spad reading and concentrations of total extracted chlorophyll in citrus leaves has been documented¹⁶. Reports are also available describing the positive linear relationship between spad readings and leaf N concentrations from different crops¹⁷⁻¹⁸. Aguilera-Gomez *et al.*¹⁹ (1999) reported that low P depresses net photosynthetic rate, stomatal conductance, P use efficiency and internal CO₂ concentration; and AM could improve all these physiological variables through improving P nutrition.

Mycorrhizal development

Interaction effect of AM inoculation and rate of P application on root colonization and spore population in rhizosphere soils of *A. mangium* was found significant (Table 3). Mycorrhizal colonization in the roots of inoculated plants given 0 and 50 g GPR per plant (63 and 61% respectively) was identical and superior to that with 100 g GPR per plant (47%). Un-inoculated plants also had some root colonization (12 to 18%) with the native AM fungi, which was much lower than that of AM inoculated plants. Spore population in the rhizosphere soils also followed similar trend as of root colonization with different rates of P application and AM inoculation.

Table 3. Interaction effect of AM inoculation and P rate on root colonization and spore population in rhizosphere soils of *Acacia mangium* on degraded land

P dose (g GPR per plant)	Root colonization (%)		Spore (No. per 10 g soil)	
	With AM	Without AM	With AM	Without AM
0	63 ^a	18 ^c	58 ^a	19 ^c
50	61 ^a	18 ^c	56 ^a	21 ^c
100	47 ^b	12 ^c	37 ^b	15 ^c
F-test	**		**	

In a column, the figures having common letter/s do not differ significantly at 1 and 5% level of probability. GPR = Gapspha phosphate rock

Results from this study is in consistent with the well-known fact that high P levels depress AM colonization while low P levels enhance AM colonization and enhance growth and nutrition of plants accordingly²⁰⁻²¹. Ammijee *et al.*²² suggest that the mechanism by which AM colonization is reduced could be related to anatomical changes of the root, making it resistant to fungal penetration by high P concentration. They also suggested that the decline in response to AM colonization is due to a combination in reduction of AM colonization as soil P increases, and the decline in the response of the plant to added P.

Tissue nutrient content

Effect of AM inoculation was significant on N and P content but not on K, Ca and Mg content (Table 4). The inoculated plants had significantly higher leaf tissue N and P content (3.90 and 0.43% respectively) compared to un-inoculated plants (3.38 and 0.38% respectively). Though leaf tissue content of K, Ca and Mg was

not significant, the inoculated plants had higher concentrations of these nutrients compared to the inoculated plants. Effect of rate of P application was significant only on leaf tissue P content (Table 4). Leaf tissue P content of plants given 50 and 100 g GPR per plant (0.41 and 0.44% respectively) was identical and significantly higher than those of control plants (0.35%). Leaf tissue content of N, K and Ca, though not significant, increased consistently with gradual increase in the rate of P application from 0 to 100 g GPR per plant. But Mg content did not follow any trend with different rates of P application. Interaction effect of AM inoculation and rate of P application on leaf tissue nutrient content was not found significant.

Table 4. Effect of arbuscular mycorrhiza (AM) and P dose on macro-nutrient content of *Acacia mangium* leaves

Treatment	Nutrient content (%)				
	N	P	K	Ca	Mg
Mycorrhiza (M)					
With M	3.90	0.43	1.02	0.66	0.22
Without M	3.38	0.38	0.83	0.59	0.21
F-test	*	*	NS	NS	NS
P dose (g GPR per plant)					
0	3.43	0.35	0.78	0.54	0.22
50	3.65	0.41	0.97	0.66	0.21
100	3.84	0.44	1.02	0.67	0.22
LSD (0.05)	NS	0.04	NS	NS	NS

NS = Not significant; GPR = Gapsha phosphate rock

Lower growth of mycorrhizal plants usually under high P fertilization and reduced photosynthetic conditions has been reported by several authors^{7,20}. Abbott and Robson²³ suggested competition between endophyte and the host plant for carbohydrates as the possible reasons for such yield depression. Callow *et al.*²⁴ stated that when P is not limiting for plant growth, (and perhaps when it is in excess), nor P will be available for carbohydrate phosphorylation, which is fundamental for active carbon transport from the host to the endophyte. Therefore, at higher soil P levels, active carbon transport can possibly proceed at higher rates, resulting in greater carbon drain from the host.

Potassium nutrition is also directly related to carbohydrate production, transport and utilization. Potassium content of inoculated plants in the current study increased steadily with increase in P level up to 100 g GPR per plant. Perhaps the combination of high tissue P and K concentration can interfere with the carbohydrate metabolism in a way that reduces carbohydrate utilization by the host plant, while allowing more carbon to be transported to the endophyte²⁵. Higher P dose given to *A. mangium* plants, increased P and K concentrations possibly to a point at which there is an inhibition on the growth promoting ability of the AM fungi, resulting in growth depression²⁵.

From the results it is evident that growth and development of *A. mangium* plants grown on degraded ex-tin mining land could

be enhanced significantly with AM inoculation. The highest growth of un-inoculated *A. mangium* plants was observed with 100 g GPR per plant; whereas for AM inoculated *A. mangium* plants the highest growth was observed with 50 g GPR per plant. This indicates that P requirement of newly planted *A. mangium* plants on degraded ex-tin mining land could be reduced to 50 g GPR per plant, *i.e.*, by 50% with AM inoculation.

References

- Lopez-Bucio J, de la Vega OM, Guevara-Garcia A & Herrera-Estrella L. 2000. Enhanced phosphorus uptake in transgenic tobacco plants that overproduce citrate. *Nature Biotechnol.* **18**: 450-453.
- Zaharah AR, Zulkifli H & Sharifuddin HAH. 1997. Evaluating the efficiency of different phosphate fertilizer sources for oil palm seedlings. *Nutr Cycl Agroecosyst.* **47**: 93-98.
- Guissou T, Ba AM, Guinko S, Duponnois R & Plenchette C. 1999. Rock phosphate and vesicular-arbuscular mycorrhiza effects on growth and mineral nutrition of *Zizyphus mauritiana* Lam in an alkaline soil. *Ann Sci For.* **55**: 925-931.
- Danneberg G, Latus C, Zimmer W, Hundes Hagen B, Schneider-Poetsch H & Bothe H. 1992. Influence of vesicular-arbuscular mycorrhiza on phytohormone balances in maize (*Zea mays* L). *J Plant Physiol.* **141**: 33-39.
- De la Cruz RE, Manalo MQ, Aggangan NS & Tanbalo JD. 1988. Growth of three legume trees inoculated with VA mycorrhizal fungi and *Rhizobium*. *Plant Soil.* **108**: 111-115.
- Marschner H & Dell B. 1994. Nutrient uptake in mycorrhizal symbiosis. *Plant Soil.* **159**: 89-102.
- Fries LLM, Pakovsky RS, Safir GR & Kaminski J. 1998. Phosphorus effect on phosphatase activity in endomycorrhizal maize. *Physiol Plant.* **103**: 162-171.
- Graham JH & Eissenstat DM. 1998. Field evidence for the carbon cost of citrus mycorrhizas. *New Phytol.* **140**: 103-110.
- Satter MA. 2000. The arbuscular mycorrhiza and phosphate rock in rehabilitation of tin tailings with *Acacia mangium* and peanut agroforestry system. *PhD Thesis*. Faculty of Agriculture, Universiti Putra Malaysia, Serdang, Malaysia.
- Koske RE & Gemma JN. 1989. A modified procedure for staining roots and detect VA mycorrhizas. *Mycol Res.* **92**: 486-505.
- Gerdemann JW & Nicolson TH. 1963. Spores of mycorrhizal endogone extracted from soil by wet sieving and decanting. *Trans Br Mycol Soc.* **46**: 235-244.
- Michigan State University. 1989. *User's Guide to MSTAT-C*. Michigan State University, East Lansing, Michigan.
- Setua GC, Kar R, Satpathy B, Das NK, Ghosh JK & Saratchandra B. 1999. Effect of vesicular arbuscular mycorrhiza on growth, leaf yield and phosphorus uptake in mulberry (*Morus alba*) under irrigated, alluvial soil conditions. *Indian J Agric Sci.* **69**: 833-836.
- Rajan SK, Reddy BJD & Bagyaraj DJ. 1999. Screening of arbuscular mycorrhizal fungi for their symbiotic efficiency with *Tectona grandis*. *For Ecol Manage.* **126**: 91-95.
- Sarkar RK, Saha A, Yamagishi Y & Saha A. 1998. Leaf positional changes in the rate of photosynthesis and specific leaf weight, chlorophyll, nitrogen content and their relationship in rice. *Indian J Plant Physiol.* **3**: 135-139.
- Duce P, Arca B, Spano D, Ventura A & Usai I. 1997. A non-destructive instrument to determine chlorophyll content: applicability to citrus. *Italus Hortus.* **4**: 26-31.
- Wu FB, Wu LH & Xu FH. 1998. Chlorophyll meter to predict nitrogen sidedress requirements for short-season cotton (*Gossypium hirsutum* L). *Field Crops Res.* **56**: 309-314.

18. Shapiro CA. 1999. Using a chlorophyll meter to manage nitrogen applications to corn with high nitrate irrigation water. *Commun Soil Sci Plant Anal.* **30**: 7-8.
19. Aguilera-Gomez L, Davies FT, Olalde-Portugal V, Duray SA & Phavaphutanon L. 1999. Influence of phosphorus and endomycorrhiza (*Glomus intraradices*) on gas exchange and plant growth of chile ancho pepper (*Capsicum annum* L cv San Luis). *Photosynthetica.* **36**: 441-449.
20. Siqueira JO, Saggin Junior OJ, Flores Aylas WW & Guimaraes PTG. 1998. Arbuscular mycorrhizal inoculation and superphosphate application influence plant development and yield of coffee in Brazil. *Mycorrhiza.* **7**: 293-300.
21. Olsen JK, Schaefer JT, Edwards DG, Hunter MN, Galea VJ & Muller LM. 1999. Effects of a network of mycorrhizae on capsicum (*Capsicum annum* L) grown in the field of five rates of applied phosphorus. *Aust J Agric Res.* **50**: 239-252.
22. Ammijee F, Stribley DP & Tinker PB. 1990. Soluble carbohydrates in the roots of leek (*Allium porrum*) plants in relation to phosphorus supply and VA mycorrhiza. *Plant Soil.* **124**: 195-198.
23. Abbott LK & Robson AD. 1986. The effect of mycorrhiza on plant growth. In *VA Mycorrhiza* (Powell CLL & Bagyaraj DJ eds), 2nd edn, pp 113-130. CRC Press, Boca Raton, Florida.
24. Callow JA, Capacio LCM, Parish G & Tinker PB. 1978. Detection and estimation of photosynthate in vesicular arbuscular mycorrhizas. *New Phytol.* **80**: 125-134.
25. Antunes V & Cardoso EJBN. 1991. Growth and nutrient status of citrus plants as affected by mycorrhiza and phosphorus application. *Plant Soil.* **131**: 11-19.

Original Article

Antimicrobial Activity of *Cinnamomum tamala* Essential Oil and Its Composition and Toxicity on White Strain Rats

Jaripa Begum^{1*}, Shoma Dutta¹, Mohammad Yusuf¹, Jashim Uddin Chowdhury¹, Kabir Ahmed¹, Salauddin Ahmed² and M Nural Anwar²

¹Bangladesh Council of Scientific & Industrial Research (BCSIR) Laboratories, Chittagong, Chittagong 4220, Bangladesh, ²Department of Microbiology, University of Chittagong, Chittagong 4331, Bangladesh

[Received 09 October 2006; Accepted 21 April 2007]

The antimicrobial activity of the essential oil from the leaves of *Cinnamomum tamala* grown at Hathazari, Chittagong was evaluated with pathogenic bacteria and phytopathogenic fungi. The aim of this study was also determine the composition of the oil and to estimate medium lethal doses (LD₅₀) of the oil after oral and intraperitoneal administration in mice. The oil was found to be very effective against all Gram-positive and Gram-negative bacterial strains tested even at very low concentration (2 µl/disk). The highest zone of inhibition was recorded with *Bacillus subtilis*, followed by *Shigella sonnei* (26 mm) at a concentration of 20 µl/disc. The oil also exhibited potent antifungal activity. The minimum concentration for absolute inhibition (MCAI) of *Fusarium equiseti* was found to be 250 ppm, followed by *Colletotrichum corchori* and *Alternaria alternata* (500 ppm). *Botryodiplodia theobromae* and *Drechslera oryzae* showed moderate sensitivity (MCAI 750 ppm) to the oil, while *Curvularia lunata* was found to least sensitive (MCAI 1,250 ppm). The leaf oil was analyzed by gas chromatograph-mass spectrometer (GC-MS). It contained 67 components representing 99.45% of the total oil constituents. The major components were eugenol (39.57%), gamma-terpinene (12.25%) and patchouli oil (10.73%). Acute toxicity test of the essential oil had been carried out on white strain rats. The oil administered orally at the rate of 2 ml/kg body weight showed no gross toxicity in rats and LD₅₀ was found to be 7.94 mg/kg body weight based on intraperitoneal administration. The essential oil of the plant presented here showed very promising activity in the area of antimicrobial agents, warranting further investigation.

Keywords: Essential oil, *Cinnamomum tamala* leaves, Antimicrobial activity, Minimum concentration for absolute inhibition (MCAI), Oil composition, Toxicity, Lethal doses 50 (LD₅₀)

Introduction

The use of plant extracts and phytochemicals, both with known antimicrobial properties, can be of great significance in therapeutic treatments. In the last few decades, a number of studies have been conducted in different countries to prove such efficiency¹. Many plants have been used because of their antimicrobial traits, which are due to compounds synthesized in the secondary metabolism of the plant. These products are known by their active substances, for example, the phenolic compounds which are part of the essential oils², as well as in tannin³. Essential oils are responsible for the fragrance of plants. These oils are secondary metabolites that are highly enriched in compounds based on an isoprene structure¹. They are called terpenes and when the compounds contain additional elements, usually oxygen, they are termed terpenoids. Terpenes or terpenoids are active against bacteria⁴⁻⁵, fungi⁶⁻⁷, viruses⁸, and protozoa⁹.

Cinnamomum tamala Nees (Lauraceae) leave, also called Bay leaf or Tejapata, is a spice of commerce. The plant is an evergreen, aromatic tree distributed in tropical and subtropical Himalayas

and in north-eastern districts of Bangladesh. Besides flavouring agent, the leaves are reported to be stimulant, carminative, diuretic, diaphoretic, lactagogue, and hypoglycaemic used in the treatment of colic, diarrhoea, anorexia, skin diseases, sore throat, coughs, colds and in scorpion sting¹⁰⁻¹². Search of literature reveals that many essential oils have good fungi-toxicity and bactericidal activity¹³⁻¹⁵. Essential oil of *C. tamala* leaves native to different geographical locations has been investigated for its composition¹⁶⁻¹⁹. However, in Bangladesh there is little information available on the composition of leaf oil of *C. tamala*²⁰ and no information available on its antimicrobial properties.

The objective of this research was to evaluate the potential of essential oil of *C. tamala* leaves on pathogenic Gram-positive and Gram-negative bacteria as well as common phytopathogenic fungi. Moreover, we investigated the toxicity and composition of the essential oil.

Material and Methods

Extraction of oil

The leaves of *Cinnamomum tamala* were collected from a single tree from the Regional Agricultural Research Station, Hathazari,

*Corresponding author:

Dr. Jaripa Begum, Principle Scientific Officer, Bangladesh Council of Scientific & Industrial Research (BCSIR) Laboratories, Chittagong, Chittagong 4220, Bangladesh
Tel (Office): (031) 681763, (031) 681763; Fax: +880 (031) 682505; E-mail: ctglab@spnctg.com

Chittagong. Voucher specimen was deposited in the herbarium of BCSIR Laboratories, Chittagong. Fresh leaves were subjected to hydrodistillation in Clevenger apparatus²¹. The oil thus obtained was dried over anhydrous sodium sulphate and stored in sealed glass vials under refrigeration prior to analysis. Refractive index of the oil was measured according to standard method²².

Test microorganisms

The Gram-positive and Gram-negative bacterial cultures used as test organisms were collected from Bangladesh Type Culture Collection (BTCC), Institute of Nutrition and Food Science, University of Dhaka and Centre for Health & Population Research, International Centre for Diarrhoeal Disease Research, Bangladesh (ICDDR,B), Mohakhali, Dhaka. Bacterial strains included *Bacillus cereus* BTCC 19, *Bacillus subtilis* BTCC 17, *Staphylococcus aureus* BTCC 43, *Bacillus megaterium* BTCC 18, *Escherichia coli* BTCC 12, *Vibrio cholerae* ICDDR, B (CRL), *Salmonella typhi* BTCC 97(CRL), *Pseudomonas sp.* BTCC 11 (CRL), *Shigella dysenteriae* BTCC 500 (CRL) and *Shigella sonnei* ICDDR,B (CRL). Phytopathogenic fungi used in this study were collected from Department of Microbiology, University of Chittagong, Chittagong. These were *Alternaria alternate*, *Curvularia lunata*, *Fusarium equiseti*, *Colletotrichum corchori*, *Drechslera oryzae*, *Macrophomina phaseolina* and *Botryodiplodia theobromae*.

Antibacterial activity assay

In vitro antibacterial activity of the essential oil was done by disc diffusion method²³ using paper disc (8 mm in diameter) and nutrient agar as basal medium. Antibacterial activities were indicated by clear zone of growth inhibition around the disc. The inhibition zones were recorded after 24 to 48 h of incubation at $37 \pm 1^\circ\text{C}$. Ampicillin (20 µg/disc) was used as standard antibiotic for comparison of results under identical condition.

Antifungal activity assay

In vitro antifungal activity of the essential oil was done by poison food technique²⁴ using potato dextrose agar (PDA) as basal medium. The diameter of radial mycelial growth of the test fungi was measured after 3 to 5 days of incubation at $25 \pm 1^\circ\text{C}$ and expressed as percent mycelial growth inhibition following the formula: $I = (C - T)/C \times 100$; where, I = percentage of inhibition, C = diameter of the fungal colony in control, T = diameter of the fungal colony in treatment. Nystatin (100 ppm) was used as standard antifungal antibiotic for comparison of results under identical condition.

Gas chromatograph-mass spectrometer (GC-MS) analysis

The analysis was carried out by GC-MS electron impact ionization (EI) method on GC-17 gas chromatograph (Shimadzu, Japan) coupled to a GCMS-QP5050A mass spectrometer (Shimadzu, Japan), fused silica capillary column, length 30 m, internal diameter (ID) 0.25 mm, inner surface coated with DB-1 (J&W), film thickness 0.25 µm, column temperature 40°C to 250°C at the rate of $4^\circ\text{C}/\text{min}$, carrier gas, helium at constant pressure of 50 kPa, acquisition parameter full scan, scan range 40-350 amu. The compounds were identified using the NIST 127 and NIST 147 library data.

Acute toxicity test on rats

For oral toxicity, the essential oil was dosed orally at the rate of 2 ml/kg body weight on 10 white strain rats (5 male and 5 female) weighing about 150-165 g/rat and observed for 24 h for any mortality and next 10 days for any delayed effect. For intraperitoneal toxicity, 60 rats weighing about 150-165 g/rat were divided into six groups. Each group received different doses of essential oil by intraperitoneal (IP) route. After the administration of different doses of essential oil, the animals were kept under observation for 24 h. The time of death were recorded. Survived animals were observed for the 0% mortality and 100% mortality was recorded according to the formulas of Miller and Tainter²⁵. The probit values were read from the table of probits. A graph of log dose versus probit was plotted. The dose corresponding to 50% or probit of 5 was recorded as lethal dose 50 (LD₅₀).

Results and Discussion

The leaf oil from *Cinnamomum tamala*, obtained with 1.0% yield value (v/w in FWB) having a refractive index of 1.468 (29°C), was a light yellow mobile liquid and possessed a spicy smell. The results of the *in vitro* antibacterial activity of the essential oil (2, 5, 10, 15 and 20 µl/disk) against the Gram-positive and Gram-negative test bacterial strains are summarized in Table 1. It was observed that the essential oil was most effective against the test bacteria strains even at the lowest concentration (2 µl/disk) of the essential oil. The highest zone of inhibition (32 mm) was recorded against *B. subtilis*; followed by *S. sonnei* (26 mm) at a concentration of 20 µl/disk. In some cases, the essential oil exhibited better antibacterial activity than the standard antibiotic, ampicillin. Similar antibacterial properties of a number of essential oils have been reported by different workers^{13,15,26-27}.

Table 1. Antibacterial activity of essential oil of *Cinnamomum tamala* leaves

Bacterium	Antibacterial activity (Diameter of zone of inhibition in mm)					
	Essential oil (µl/disk)					Ampicillin
	2	5	10	15	20	(20 µg/disk)
Gram-positive organisms						
<i>Bacillus cereus</i>	18	19	21	22	24	22
<i>Bacillus megaterium</i>	17	19	21	22	24	22
<i>Bacillus subtilis</i>	20	24	25	28	32	25
<i>Staphylococcus aureus</i>	15	16	18	20	21	20
Gram-negative organisms						
<i>Escherichia coli</i>	14	17	18	21	23	13
<i>Pseudomonas</i> species	15	20	21	22	24	19
<i>Salmonella typhi</i>	16	19	20	21	22	30
<i>Shigella dysenteriae</i>	16	20	21	22	24	35
<i>Shigella sonnei</i>	18	21	24	25	26	30
<i>Vibrio cholerae</i>	15	19	20	21	23	24

The *in vitro* antifungal activity of the essential oil and standard antibiotic nystatin is shown in Table 2. It was revealed from the results that the antifungal activity of the oil was remarkable; inhibiting the growth of all the 7 test fungal pathogens completely (100%). The minimum concentration for absolute inhibition (MCAI) of the oil against *F. equiseti* was found to be 250 ppm; followed by *C. corchori* and *A. alternata* (500 ppm). *B. theobromae*, *D. oryzae* and *M. phaseolina* showed more resistance towards the oil and MCAI was found to be 750 ppm; followed by *C. lunata* (1,250 ppm). At lower concentration, the oil exhibited fungistatic activity, but at MCAI it was fungicidal. It was also observed that the essential oil was more effective against the phytopathogenic fungi than those of the bacterial strains. Similar antifungal properties of a number of essential oils have been reported by several workers²⁸⁻³².

Table 2. Antifungal activity of essential oil of *Cinnamomum tamala* leaves

Bacterium	Antifungal activity (Percent growth inhibition)						
	Essential oil (ppm)						Nystatin (100 ppm)
	100	250	500	750	1,000	1,250	
<i>Alternaria alternata</i>	50	78	100	100	100	100	56
<i>Botryodiplodia theobromae</i>	13	33	80	100	100	100	82
<i>Colletotrichum corchori</i>	50	78	100	100	100	100	42
<i>Curvularia lunata</i>	27	56	72	80	88	100	72
<i>Drechslera oryzae</i>	7	9	32	100	100	100	76
<i>Fusarium equiseti</i>	29	100	100	100	100	100	46
<i>Macrophomina phaseolina</i>	25	33	52	100	100	100	71

The essential oil was analyzed by GC-MS and many components were identified. Some of components that might have antimicrobial properties are listed in Table 3. Sixty seven components representing 99.45% of the total leaf oil were identified. Eugenol (39.57%), gamma terpinene (12.25%) and patchouli oil (0.73%) were the major components in the leaf oil. Eugenol is a well-characterized representative of phenolic compounds possessing a C₃ side chain at a lower level of oxidation and containing no oxygen are classified as essential oils¹. Eugenol is considered bacteriostatic against both fungi³³ and bacteria³⁴.

The acute toxicity test of the leaf essential oil was carried out on white strain rats, orally and intraperitoneally. The oil administered orally at the rate of 2 ml/kg body weight showed no gross toxicity in rats. The results of intraperitoneal toxicity test are presented in Table 4. The LD₅₀ value was found to be 794 mg/kg body weight. These results suggest the presence of bioactive plant metabolites.

Table 3. Compositions of the essential oil of *Cinnamomum tamala* leaves

Component	Percentage
Bornylene	0.06
3-Carene	2.07
Bicyclo [3.1.0] hex-2-ene, 4-methyl-1-(1-methyl-1-yl)-	3.26
Fenchene	0.19
Camphene	0.16
Sabinene	0.15
Beta-pinene	2.40
Gamma-terpinene	12.25*
Cyclobutane, 1,2-bis (1-methylethenyl)-, trans-	5.77
Eucalyptol	2.44
1,3,6-Octatriene,3,7-dimethyl-,(Z)	1.01
alpha-Thujene	0.18
1,4-Cyclohexadiene,1-methyl-4-(1-methylethyl)-	0.56
Cyclohexene,1-methyl-4-(1-methylethyl)-	0.36
alpha-Terpinene	2.84
Isocrotylbenzene	0.05
2-Cyclohexen-1-ol,1-methyl-4-(1-methylethyl)-, cis-	0.16
Cycloheptane, 1,3,5-tris(methylene)-	0.04
Borneol	0.39
L-4-terpineneol	0.42
p-Cymen-8-ol	0.81
Terpinyl acetate	1.95
Sabinyl acetate	0.48
cis-Piperitol	0.04
Linalool	0.17
p-Cumenol	0.07
Lavaandulol	0.18
Esdragole	0.05
Thymol	0.04
Bornyl acetate	0.09
Dihydroedulan IIA	0.12
Elixwne	0.13
Eugenol	33.57*
Caryophyllene	0.16
alpha-Guaiene	0.62
Ethylmethyl imidazole	0.14
alpha-Panansin	0.40
alpha-Patchoulene	0.33
Patchoulene	0.09
Germacrene D	0.18
Eugenyl acetate	6.00
Decahydro-8a-ethyl-2,1,4a,6-tetramethylnaphthalene	0.06
beta-Elementene	0.09
Spiro [androst-5-ene-17, 1p-cyclobutan]	
2p-one, 3-hydroxy-, (3.beta., 17.beta.)	0.06
Diepi-alpha-cedrene epoxide	2.33
Caryophylleno oxide	0.67
alpha-Guaiene	0.13
Spathulenol	0.67
Viridiflorol	0.91
Cubenol	0.22
Ledol	1.49
Patchouli alcohol	10.37*
Carotol	0.14
alpha-Bisabolol	0.21
Tetrahydroedulan	0.93
trans, trans-Farnesal	0.26
beta-Humulene	0.14
3-Nonen-5-one	0.05
delta.-Neoclovene	0.04
cis-(2,4a,5,6,9-tetramethyl (1H) benzocycloheptene	0.10
3,7,11-Trimethyl-dideca-2, 4,6,10-tetraenal	0.13
2(1H) Naphthalenone, 3,5,6,8,	
8a-hexahydro-4,8a-dimethyl-6-(1-methylethenyl)-	0.05
Humulen (VI)	0.08
1-Cyclohexene-1-propanal, 2,6,6-trimethyl-	0.09
Corymbolone	0.15
Pregnan-20-one,3-(acetyloxy)-e-hydroxy-6,	
16-dimethyl-,(3. beta., 5. alpha., 6.beta., 16. alpha.)-	0.07

*Major components

Table 4. Toxicity test of the essential oil of *Cinnamomum tamala* leaves based on intraperitoneal (IP) administration to white strain rats

Rat group	Dose (mg/kg)	Concentration (ml/rat)	Dead (No.)	Survived (No.)	Death rate (%)	Corrected (%)	Probit
1	1,800	0.25	10	0	100	97.5	6.97
2	1,440	0.20	8	2	80	80.0	5.84
3	1,080	0.15	6	4	60	60.0	5.25
4	720	0.10	4	6	40	40.0	4.75
5	360	0.05	2	8	20	20.0	4.61
6	180	0.026	0	10	0	2.5	3.04

The antimicrobial properties of plants have been investigated by a number of researchers worldwide. It was documented that several compounds extracted from these plants inhibited the growth of Gram-positive and Gram-negative organisms and some of them also inhibited the growth of fungi³⁵⁻³⁸. More detailed study on antimicrobial compounds was done evaluating extracts from 120 plant species from 28 different families³⁹. It was documented that 81 extracts obtained from 58 plants were active against *S. aureus*, and five extracts from four other plants inhibited the growth of *P. aeruginosa*. Another study⁴⁰ detected the antibacterial and antifungal (*C. albicans*) activity of essential oils obtained from *Croton triangularis* leaves.

Since the products of higher plant origin are reported to be biodegradable, the *C. tamala* leaf oil with its low toxicity, strong antifungal and antibacterial properties could be an excellent candidate as a natural antimicrobial agent. Although the active components in the extracts responsible for antibacterial and antifungal activities have not been identified conclusively, all of these results imply the existence of bioactive metabolites and therefore, more research with this plant is warranted.

References

- Cowan MM. Plant products as antimicrobial agents. *Clin Microbiol Rev.* **12**(4): 564-582.
- Jansen AM, Cheffer JJC & Svendsen AB. 1987. Antimicrobial activity of essential oils: A 1976-1986 literature review. Aspects of test methods. *Planta Med.* **40**: 395-398.
- Saxena G, McCutcheon AR, Farmer S, Towers GHN & Hancock REW. 1994. Antimicrobial constituents of *Rhus glabra*. *J Ethnopharmacol.* **42**: 95-99.
- Barre JT, Bowden BF, Coll JC, Jesus J, Fuente VE, Janairo GC & Ragasa CY. 1997. A bioactive triterpene from *Lantana camara*. *Phytochemistry.* **45**: 321-324.
- Habtemariam S, Gray AI & Waterman PG. 1993. A new antibacterial sesquiterpene from *Premna oligotricha*. *J Nat Prod.* **56**: 140-143.
- Harrigan GG, Ahmad A, Baj N, Glass TE, Gunatilaka AAL & Kingston DGI. 1993. Bioactive and other sesquiterpenoids from *Porella cordeana*. *J Nat Prod.* **56**: 921-925.
- Rana BK, Singh UP & Taneja V. 1997. Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. *J Ethnopharmacol.* **57**: 29-34.
- Hasegawa H, Matsumiya S, Uchiyama M, Kurokawa T, Inouye Y, Kasai R, Ishibashi S & Yamasaki K. 1994. Inhibitory effect of some triterpenoid saponins on glucose transport in tumor cells and its application to *in vitro* cytotoxic and antiviral activities. *Planta Med.* **6**: 240-243.
- Ghoshal S, Prasad BNK & Lakshmi V. 1996. Antiamoebic activity of *Piper longum* fruits against *Entamoeba histolytica* *in vitro* and *in vivo*. *J Ethnopharmacol.* **50**: 167-170.
- Chopra RN, Nayer SL & Chopra IC. 1956. *Glossary of Indian Medicinal Plants*, pp 65-66. Council of Scientific and Industrial Research, New Delhi.
- Yusuf M, Chowdhury JU, Wahab MA & Begum J. 1994. *Medicinal Plants of Bangladesh*, p 63. Bangladesh Council of Scientific and Industrial Research, Dhaka.
- Ghani A. 1998. *Medicinal Plants of Bangladesh: Chemical Constituents*, 1st edn, p 128. Asiatic Society of Bangladesh, Dhaka.
- Maruzella JC & Henry PA. 1958. The antibacterial activity of perfume oils. **47**: 471-476.
- Sinha GK & Gulati BC. 1990. Antibacterial and antifungal study of some essential oils and some of their constituents *Indian Perfumer.* **34**(2): 126-129.
- Begum J, Chowdhury JU, Yusuf M, Wahab MA, Ahmed K, Akter N & Anwar MN. 1999. Antimicrobial activity of essential oils isolated from the *Ocimum amricanum* and *O. gratissimum* var. *clocimum* and their toxicity in white strain rats. *Bangladesh J Microbiol.* **16**(2): 127-134.
- Sood SP, Padha CD, Talwar YP, Jamwal RK, Chopra MM & Rao PR. 1979. Essential oils from the leaves of *Cinnamomum tamala* Nees & Eberm growing in Himachal Pradesh. *Indian Perfumer.* **23**: 75-78.
- Gulati BC. 1982. Essential oils of *Cinnamomum* species. In *Cultivation and Utilization of Aromatic Plants* (Atal CK & Kapur BM eds), pp 607-619. Regional Research Laboratory (CSIR), Jammu-Tawi.
- Nath SC, Singh RS & Hazarika AK. 1994. Essential oil of leaves of *Cinnamomum tamala* Nees & Eberm from North East India. *J Spices Aromatic Crops.* **3**: 33-35.
- Showkat RM, Ali M & Kapoor R. 2004. Chemical composition of essential oil of *Cinnamomum tamala* Nees & Eberm leaves. *Flav Frag J.* **19**: 112-114.
- Begum J, Chowdhury JU, Yusuf M & Wahab MA. 1993. Studies on essential oils for their antibacterial and antifungal properties. Part 1. Preliminary screening of 35 essential oils. *Bangladesh J Sci Ind Res.* **28**(4): 25-34.
- Clevenger JF. 1928. Apparatus for the determination of volatile oil. *J Am Pharm Assoc.* **17**: 346.
- Whitely MA. 1947. *Thorpe's Dictionary of Applied Chemistry*, 4th edn, Vol 8, p 651. Green and Co Ltd, Longmans.
- Bauer AW, Kibry MM, Sherris JC & Turck M. 1966. Antibiotic susceptibility testing by a standardized single disc method. *Am J Clin Path.* **45**: 493-496.
- Grover RK & Moore JD. 1962. Toximetric studies of fungicides against brown rot organisms *Sclerotinia fructicola* and *S. laxa*. *Phytopathol.* **52**: 876-880.

25. Miller LC & Tainter ML. 1944. Estimation of the ED₅₀ and its error by means of logarithmic-probit paper. *Proc Soc Exp Biol Med.* **59**: 261-264.
26. Rao RSS & Nigam SS. 1978. Chemical and antimicrobial examination of the essential oil of *Nigella sativa* var. *aromatica*. *Indian Perfumer.* **22**(4): 232-238.
27. Chowdhury DC, Suri RK & Deshmukh DK. 1998. *In vitro* antimicrobial activity of essential oil of newly evolved *Eucalyptus* hybrid FRI-4. *Indian Forester.* **114**(1): 35-38.
28. Thind TS & Suri KR. 1979. *In vitro* antifungal efficacy of four essential oils. *Indian Perfumer.* **23**(2): 138-140.
29. Handique AK & Singh HB. 1990. Antifungal action of lemongrass oil on some soil born plant pathogens. *Indian Perfumer.* **34**(3): 232-234.
30. Begum J, Chowdhury JU, Yusuf M, Wahab MA, Ahmed K, Akter N & Anwar MN. 1997. Evolution of oils extracted from *Eucalyptus citriodora* and *E. camaldulensis* for antimicrobial properties and their toxicity in rats. *Bangladesh J Sci Ind Res.* **32**(4): 561-566.
31. Chowdhury JU, Begum J, Yusuf M & Wahab MA, Kamal AKMM & Chowdhury SA. 1999. Studies on *Cymbopogon flexuosus* for its growth, yield performances, essential content, physicochemical characteristics and fungitoxic properties of the oil. *Bangladesh J Sci Ind Res.* **34**(3-4): 418-421.
32. Chowdhury JU, Yusuf M, Begum J, Sultana SA & Husain MM. 2003. Composition and fungitoxic properties of the essential oil of *Lippia javanica* leaves. *Indian Perfumer.* **47**(4): 385-388.
33. Duke JA. 1985. *Handbook of Medicinal Herbs*. CRC Press Inc, Boca Raton, Florida.
34. Thomson WAR. 1978. *Medicines from the Earth*. McGraw-Hill Book Co, Maidenhead.
35. Alonso-Paz E, Cerdeiras MP, Fernandez J, Ferreira F, Moyna P, Soubes M, Vazquez A, Veros S & Zunno L. 1995. Screening of Uruguayan medicinal plants for antimicrobial activity. *J Ethnopharmacol.* **45**: 67-70.
36. Anesini E & Perez C. 1993. Screening of plants used in Argentine folk medicine for antimicrobial activity. *J Ethnopharmacol.* **39**: 119-128.
37. Martinez MJ, Vasquez SM, Espinosa-Perez C, Dias M & Herrera-Sanchez M. 1994. Antimicrobial properties of argentatine - a isolated from *Parthenium argentatum*. *Fitoterapia.* **65**: 371-372.
38. Martinez MJ, Betancourt J, Alonso-Gonzalez N & Jauregui A. 1996. Screening of some Cuban medicinal plants for antimicrobial activity. *J Ethnopharmacol.* **52**: 171-174.
39. Santos Filho D, Sarti SJ, Bastos JK, Leitão Filho HF, Machado JO, Araujo MLC, Lopes WD & Abreu JE. 1990. Atividade antibacteriana de extratos vegetais. *Rev Cien Farm.* **12**: 39-46.
40. Lemos TLG, Monte FJQ, Matos FJA, Alencar JW, Craveiro AA, Barbosa RCSB & Lima ED. 1992. Chemical composition and antimicrobial activity of essential oils from Brazilian plants. *Fitoterapia.* **63**: 266-268.